STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO. 9812.2 DISC.

Decident®

Disposable Disinfectant Sleeve

U.S. Pat. No. 4,446,967

ACTIVE INGREDIENTS:

2-Phenylphenol	0.249%
4-tert-Amyphenol	0.058%
Ethanol*	66.000%
INERT INGREDIENTS:	33.693%
TOTAL	100.000%

*Specially Denatured Alcohol

KEEP OUT OF REACH OF CHILDREN

WARNING

See back panel for additional precautionary statements.

NET CONTENTS: 24 SLEEVES EACH WITH 1/2 FL. OZ. (15 ML)



Manufactured for:

NEVIN LABORATORIES INC.

EPA Reg. No. 10088-105-58520

EPA Est. No. 054450-IL-001

FEATURES, CLAIMS, AND USES

HOSPITAL-TYPE DISINFECTING ACTION: Medical-level and broad spectrum antimicrobial activity destroys many organisms which threaten public health or create unpleasant odors and other undesirable effects. Reduces the hazard of cross-contamination from hard, inanimate surfaces in medical facilities.
VIRUCIDAL*- Inactivates these agents: Human Immunodeficiency virus Type 1(HIV-1); a retrovirus that causes "AIDS" (Acquired Immune Deficiency Syndrome); Herpes simplex Type 1 (a herpes virus that causes cold sores); herpes simplex Type 2 (a herpes virus that causes "genital herpes"); Coxsackle viruses A2 and B3 (picornaviruses; human enteroviruses); Adenovirus Type 5 (an adenovirus); Influenza A2/Hong Kong (a myxovirus); and Vaccinia (a poxvirus) on hard inanimate surfaces.
BACTERICIDAL- kills Escherichia coli; Pseudomonas aeruginosa; Salmonella choleraesuis; Salmonella schottmuelleri (paratyphoid B); Shigella dysenteriae; Staphylococcus aureus; and Streptococcus pyogenes.
TUBERCULOCIDAL - Mycobacterium tuberculosis (Tubercle bacilli) on surfaces that have been thoroughly cleaned prior to treatment. Effective in 10 minutes at 20°C (68°F).

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye irritation. Do not get in eyes or on skin. Avoid breathing spray mist or vapors. Harmful if swallowed.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control treatment center or doctor for further treatment advice.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

PHYSICAL OR CHEMICAL HAZARDS

FLAMMABLE: Keep away from heat, sparks, and flames. Do not use in the vicinity of pilot lights. Do not use on latex paint, polished wood furniture, or other materials that could be damaged by alcohol.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.
TO KILL HIV-1 (Human immunodeficiency virus Type 1, AIDS virus) ON PRECLEANED SURFACES in health care or other settings with inanimate environmental surfaces/objects associated with the potential for transmission of HIV-1 and likely to be soiled with blood/body fluids, follow these ADDITIONAL SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:
PERSONAL PROTECTION-Barrier protection items such as disposable latex gloves, gowns, masks, and eye coverings should be worn when handling items soiled with blood/ body fluids. CLEANING PROCEDURE-Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product. CONTACT TIME-Leave surface wet for 10 minutes. DISPOSAL OF INFECTIOUS MATERIALS-Blood/body fluids and materials removed from or used to clean surfaces/objects should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

FOR DISINFECTING DENTAL HANDPIECES

NOTE: The disposable dental handpiece sleeve is not a substitute for sterilization. It is recommended only for disinfecting handpieces which cannot be sterilized by steam autoclave, unsaturated chemical vapor, dry heat, or ethylene oxide, due to physical limitations, incompatibility, or residue.

FOLLOW THESE STEPS!

Always wear gloves. Flush handpiece thoroughly for 20 to 30 seconds. Remove the bur. Clean handpiece thoroughly to remove saliva, blood, and other organic soil. Scrub handpiece with small brush and detergent under running water. Blot handpiece dry with a clean paper towel. Water droplets clinging to the handpiece will dilute the disinfectant. Grasp tabs at upper right corner of pouch and open to the line "open here". This allows sufficient room to insert either a high or low speed handpiece. Insert handpiece completely into sleeve between the disinfectant-soaked sponges. Vigorously massage handpiece inside sleeve from top to bottom. Massaging action distributes disinfectant over entire handpiece surface and provides a beneficial scrubbing effect. Leave handpiece inside sleeve for at least 10 minutes at approximately 20°C (68°F). Handpiece with sleeve may be suspended by tubing with open end of sleeve upward. After 10 minute disinfection cycle, rinse handpiece in water, or wipe with an alcohol-saturated sponge to remove residual disinfectant. Blot dry with a paper towel. Lubricate handpiece according to manufacturer's directions. Dispose of the used sleeve in the trash.

STORAGE AND DISPOSAL

GENERAL: Do not contaminate other materials (including water/food/feed) during transport, use, storage, and disposal. If damaged, leaking, or spilled, soak up and wrap any waste, then dispose of as below.
STORAGE: Keep unused sleeves in original carton in a cool, dry, locked area inaccessible to children and away from sources of ignition, heat, degradation, and contamination (including flames, sparks, hot surfaces, sunlight, and other chemicals).
DISPOSAL OF SLEEVE: Do not reuse sleeve; after use, discard in trash.

5000 S. Halsted Street
Chicago, IL 60609
(773) 624-4330
(800) 544-5337
FAX (773) 624-7337

Manufactured for:

NEVIN LABORATORIES INC

EPA Reg. No. 10088-105-58520
EPA Est. No. 054450-IL-001
Made in U.S.A.

25560
1066 Form 59