Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2009-2011 LIC. NO.

9822.14

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2012-2014 LIC. NO.

9822.14

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO.

9822.14

SIPCAM AGRO USA, INC.

CAVALCADE™ 65WDG

For preemergence control of grass and broadleaf weeds in:

- Established turfgrasses (excluding golf course putting greens), lawns and sod nurseries
- Landscape ornamentals in nurseries or in established plantings
- Established perennials and wildflower plantings
- Plants grown for cut foliage production (Florida only)
- Conifer and hardwood tree seedling nurseries
- Christmas tree farms
- Managed transportation and utility rights-of-way, including rail and equipment yards, and public utility facilities (substations, tank farms, pumping stations, parking/storage areas, ungrazed fencerows)

ACTIVE INGREDIENT:	
Prodiamine (CAS No. 29091-21-2)	65.0%
OTHER INGREDIENTS:	35.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have affected person sip a glass of water if able to swallow. • Do not induce vomiting unless told by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 900-4044 Poison Control Center (human health) (800) 345-4735 ASPCA (animal health)

Cavalcade is a trademark of Sipcam Agro USA, Inc.

NET CONTENTS: 10 pounds

SIPCAM AGRO USA, INC.
300 COLONIAL CENTER PARKWAY, SUITE 230
ROSWELL, GA 30076

EPA REG. NO. 60063-26
EPA EST. NO. 37429-GA-01

RECEIVED
AUG 13 2009

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact, while mixing or handling the concentrated material, may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT

WPS USES:

Mixers, loaders and applicators and other persons who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR part 170) – in general, agricultural-plant uses are covered – must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves, such as butyl rubber >14 mils, or neoprene rubber > 14 mils, or nitrile rubber >14 mils (See instructions for Category A on the EPA chemical resistance category selection chart B if you want other options.)
- Shoes plus socks

NON-WPS USES:

Mixers and loaders who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR part 170) – in general, only agricultural-plant uses are covered by the WPS – must wear:

- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- After handling this product, immediately wash the outside of gloves before removing them, then remove gloves and all other PPE. Immediately wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFF part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

(continued)

AGRICULTURAL USE REQUIREMENTS *(continued)*

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI)of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

WHERE TO USE

Cavalcade 65WDG is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turfgrasses (excluding golf course putting greens), lawns and sod nurseries
- Landscape ornamentals in nurseries or in established plantings
- Established perennials and wildflower plantings
- Plants grown for cut foliage production (Florida only)
- Conifer and hardwood tree seedling nurseries
- Christmas tree farms
- Managed transportation and utility rights-of-way, including rail and equipment yards, and public utility facilities (substations, tank farms, pumping stations, parking/storage areas, ungrazed fencerows

HOW CAVALCADE 65WDG WORKS

Cavalcade 65WDG controls susceptible weeds by preventing growth and development of newly germinated weed seeds. Weed control is most effective when Cavalcade 65WDG is activated by at least 0.5 inch of rainfall or irrigation or shallow incorporation (1 to 2 inches) before weed seeds germinate and within 14 days following application.

USE PRECAUTIONS

1. Do not graze or feed livestock forage cut from areas treated with Cavalcade 65WDG.
2. Follow all applicable directions, restrictions, and precautions on the labels of EPA-registered tank mix partners.
3. Do not blend Cavalcade 65WDG onto dry fertilizer or any other granular material.
4. Chemigation: Do not apply this product through any type of irrigation system unless instructed otherwise in this label.
5. Do not apply aerially.
6. Do not apply to golf course putting greens.

MIXING AND APPLICATION

Mixing

Cavalcade 65WDG must be mixed thoroughly in the spray tank to ensure uniform application. Follow these steps:

1. Fill the spray tank ¼ full with clean water or fluid fertilizer only.
2. Start agitation and check to ensure it is working properly.
3. Add Cavalcade 65WDG directly into the tank.
4. Add the rest of the carrier to obtain the final spray volume.
5. A spray colorant may be used with Cavalcade 65WDG to mark areas as they are treated. This will improve application accuracy by minimizing swath skips and overlaps.
6. Maintain vigorous agitation in the spray tank before and during the application. This will ensure a well-mixed spray suspension.
7. Do not allow spray suspension to dry in the tank. Thoroughly clean the sprayer after use by flushing the system with water containing a detergent. Refer to the Pesticide Disposal section of this label for waste disposal.

Tank Mixing Cavalcade 65WDG

Cavalcade 65WDG may be tank mixed with certain other EPA-registered herbicides to provide a broader spectrum of weed control or to control emerged weeds. Refer to the specific directions for use for tank mix partners, and consult the label(s) of the individual tank mix partners(s) for use rate, application timing, weeds controlled, and specific precautions and/or restrictions. Tank mixes are permitted only in states where the tank mix partners(s) are registered for the application site and the turf and ornamental species listed. When using Cavalcade 65WDG in a tank mixture with other pesticides, observe the most restrictive label limitations and precautions on the labels of the products used.

Before tank mixing pesticides, it is advisable to test compatibility by mixing the products in a small container first. See the Compatibility Test section.

Compatibility Test

Before mixing Cavalcade 65WDG with other pesticides in the spray tank, test the compatibility by mixing all components (carrier and pesticide products) in a small container in proportionate quantities. For example, a 1-qt. Jar would be 1/100 the volume of a 25 gals./A spray rate. At 1 lb./A the Cavalcade 65WDG rate would be proportional to 4.5 g per qt. Add approximately 1.5 teaspoons to a qt. of water. Calculate amounts for the other products based on rate per acre. An approximate volume would be 1.5 teaspoons for each lb./A of a dry formulation and 0.5 teaspoons for each pt./A of a liquid formulation. (See following table.)

Amount of Component to Add to One quart Jar of Spray Carrier (Assuming Carrier Volume of 25 gals./A)

Component Formulations	Rate Per		Level Teaspoons
	Acre	1,000 sq. ft.	
Cavalcade 65WDG	1.0 lb.	0.4 oz.	1.5
Dry Tank Mix Partners	1.0 lb.	0.4 oz.	1.5
Liquid Tank Mix Partners	1.0 pt.	0.4 oz.	0.5

If components do not ball-up or form flakes, sludge, gels, oily films or layers, then the mixture is compatible. Incompatibility will usually occur within 5 minutes after mixing. If the components are not compatible, use a compatibility agent and rerun the test to determine if the mixture is suitable. If components are still not compatible, do not tank mix.

Mixing Order for Tank Mixtures

Notes: (1) When mixing Cavalcade 65WDG with other components (carrier and partner pesticide products), allow products to completely dissolve between steps. This is key when tank mixing with ester formulations. (2) Maintain agitation throughout mixing and application of the mixture.

Add the products to the spray tank in the following order:

1. Add products packaged in water-soluble bags first. Agitate the tank mixture. Allow the water-soluble bags to completely dissolve and the product to disperse before adding any other tank mix partner.
2. Then add water-dispersible granules (WDG or WG formulations) and wettable powders (WP formulations). Add wettable powders to the tank as agitation continues. Allow the product to disperse completely before other products are added.
3. Add spray adjuvants and spray markers. Read the adjuvant's label first and use only those adjuvants approved for application to turf and ornamentals.
4. Add flowable liquids (FL) or suspension concentrates (SC).
5. Add emulsifiable concentrates (EC) last.

Application

Apply Cavalcade 65WDG in a minimum of 20 gals/A (0.5 gal./1,000 sq. ft.) of carrier (water and/or fluid fertilizer) using a calibrated, low-pressure sprayer with 50-mesh or coarser screens. A broadcast boom or handheld wand designed for herbicide or insecticide application will provide the best results. Select nozzle pressure and gallonage to provide complete coverage.

Weeds Controlled

When used as directed in this label, Cavalcade 65WDG will control the following weeds:

Barnyardgrass	Goosegrass [5]	Purslane, Common
Bluegrass, Annual (*Poa annua*) [1]	Henbit [2]	Pusley, Florida
Carpet weed	Itchgrass	Rescuegrass [4]
Chickweed, Common [2]	Johnsongrass (from seed)	Shepherds Purse [2]
Chickweed, Mouseear (from seed)	Junglerice	Signalgrass, Broadleaf
Crabgrass (Large, Smooth) [3]	Knotweed [2]	Speedwell, Persian
Crowfootgrass	Kochia	Sprangletop
Cupgrass, Woolly	Lambsquarter, Common	Spurge, Prostrate
Foxtails, Annual	Lovegrass	Witchgrass
Pigweed	Panicum (Texas, Fall, Browntop)	Woodsorrel, Yellow (from seed)

[1] In areas where *Poa annua* is a winter annual, apply Cavalcade 65WDG (see rate table) in August or September to established, non-overseeded turf before *Poa annua* seeds germinate. These timings are approximate. Consult State Extension Service for

more specific timing for your area. Also see the section of this label ***Poa Annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV, and TX only).**

[2] To control this weed, apply Cavalcade 65WDG in late summer, fall, or winter before weed seeds germinate.

[3] **Fall Applications for Spring Crabgrass Control in Cool-Season Grasses:** In those areas where the ground freezes in the winter, Cavalcade 65WDG can be applied in the fall at rates of 1.0-1.15 lbs./A after the soil temperature falls below 50°F but before the ground freezes. This application will control crabgrass the following spring.

[4] Suppression only.

[5] In any area a single application of 1-2.3 lbs./A of Cavalcade 65WDG will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective control may be obtained by making an initial application of 1-1.5 lbs./A followed, after 60-90 days, by a second application that does not exceed the maximum rate for that turfgrass species listed in the **Maximum Application Rate Table**.

ESTABLISHED TURF

Cavalcade 65WDG is a preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turfgrasses including:

- Golf courses **excluding** putting greens
- Lawns
- Sod nurseries

The maximum amount of Cavalcade 65WDG that may be applied per year is given for each turfgrass species in the Annual Use Rates section of the label.

For optimum weed control, Cavalcade 65WDG should be activated by at least 0.5 inch of rainfall or irrigation before weed seeds germinate and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



Use Precautions-Turfgrass

1) Do not apply Cavalcade 65WDG to areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
2) Do not cut (harvest) treated sod before 90 days after application. To avoid turfgrass injury, do not apply to newly set sod until the sod has rooted and exposed edges have filled in.
3) To avoid turfgrass injury, do not apply Cavalcade 65WDG to turf stressed by conditions such as drought, low fertility, or pest damage.
4) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
5) ***Do not apply Cavalcade 65WDG to golf course putting greens.***
6) If the depth of the creeping bentgrass root system becomes shallow and root tips contact Cavalcade 65WDG-treated soil, new root formation may be inhibited. Mowing height can affect the depth of a plant's root system. To avoid this, do not apply Cavalcade 65WDG to creeping bentgrass less than 0.5 inch in height.

Application Timing and Rate-Turfgrass

Cavalcade 65WDG may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications should be made before target weeds germinate.

Cavalcade 65WDG will not control weeds that have already emerged.

The amount of Cavalcade 65WDG to apply depends upon:

1. the length of residual weed control desired (the higher the application rate, the longer the control),
2. the turf species, and
3. the maximum amount which can be applied to the turf species per calendar year.

Length of Crabgrass Control*



***Length of control varies by region. This table is an average for planning purposes.**

Annual Use Rates-Turfgrass

Cavalcade 65WDG can be applied to the turfgrass species listed in the following table. Do not apply more than the highest rate listed for each species in a calendar year.

Table 1.
Maximum Application Rate of Cavalcade 65WDG Per Calendar Year by Turfgrass Species [1]

Turf Species	lbs. Product/Acre	oz. Product/1,000 sq. ft.
Bermuda grass [2] Bahiagrass Centipedegrass Kikuyugrass Seashore Paspalum St. Augustinegrass [3] Tall Fescue (including turf-type) Zoysiagrass	1.0-2.30 [1]	0.36-0.83
Buffalograss Kentucky Bluegrass Perennial Ryegrass	0.5-1.50 [1]	0.185-0.55
Fine Fescue	0.5-1.15 [1]	0.185-0.42
Creeping Bentgrass (0.5 inches or more in height) [4]	0.5-1.00 [1]	0.185-0.37

[1] Cavalcade 65WDG may be applied more than once a year as long as the total amount applied is not greater than the maximum application rate for each turf species. All applications must be made before weed seeds germinate.

[2] May be used on newly-sprigged or plugged Bermudagrass at rates not to exceed 0.80 lb./A (0.30 oz./1,000 sq. ft.). Newly-sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded.

[3] Use an initial rate of 0.75-1.5 lbs./A per application.

[4] To avoid grass injury, do not apply Cavalcade 65WDG to creeping bentgrass mowed at less than 0.5 inch in height.

When to Apply Cavalcade 65WDG After Overseeding Turf

Injury to desirable seedlings is likely if Cavalcade 65WDG is applied before the secondary roots of seedlings are in the second inch of soil (not thatch plus soil). To reduce the potential to injure overseeded turf, wait 60 days after seeding or until after the second mowing, whichever is longer, before applying Cavalcade 65WDG.

When to Overseed After Application-All States*

Cavalcade 65WDG will inhibit the development of turfgrass species overseeded too soon after application.

Follow rates and intervals in the table below for best overseeding/reseeding results.

*Note: In AZ, CA, NV, and TX, the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section "*Poa Annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV, and TX only)".

Amount of Cavalcade 65WDG	Interval (Months) Before Overseeding		
Lbs. Product/A	North	Transition	South
.75	4	4	4
1.00	5	4	4
1.15	6	5	5
1.25	-	6	6
1.50	-	7	7
1.75	-	-	9
2.00	-	-	10
2.30	-	-	12

Poa annua control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV, and TX only)

Use on golf courses (excluding golf course putting greens, lawns, and sod nurseries when overseeding with perennial ryegrass (minimum seeding rate of 350 lbs./A).

How Much and When to Apply

Amount to Apply	When to Apply	Expected Control	Use Precautions
0.58-1.0 lb./A	**First application:** 6 to 8 weeks **before** ryegrass overseeding **Second application:** 4 to 8 weeks **after** overseeding or when perennial ryegrass roots are in the second inch of soil	1 application for 70% or greater control of *Poa annua* Second application may enhance control	1. Some seedling mortality and temporary reduction in root growth of new seedlings may occur. 2. To reduce the potential for seedling mortality, maintain a moist seedbed with light, frequent irrigation. 3. make no more than 2 applications per year for this use, and do not exceed a total of 1.3 lbs./A per year. 4. Do not make a second application if any injury to the ryegrass is observed after the first application. 5. Do not make a second application unless the product was first applied before overseeding.

ORNAMENTALS (CONTAINER, FIELD, AND LANDSCAPE GROWN, INCLUDING CHRISTMAS TREE FARMS), RIGHTS-OF-WAY, GROUNDS OF UTILITIES, UNGRAZED FENCE ROWS

Cavalcade 65WDG may be applied to soil surfaces for preemergence control of many grass and broadleaf weeds:

- Around ornamental shrubs, trees, established perennial vegetation and wildflower plantings;
- On or surrounding managed rights-of-way for transportation systems including roadways, roadsides, railways, and equipment yards;
- On grounds of utilities such as power substations, tank farms, pumping stations, parking and storage areas;
- On ungrazed fence rows.

Application Timing and Information

Cavalcade 65WDG:

1. Will not control emerged weeds.
2. May be applied to newly-transplanted and established ornamentals as broadcast or over-the-top-spray.
3. Is most effective when applied to soil free of clods, weeds, and debris such as leaves and mulch.
4. Is most effective when the product is activated in the soil before weed seeds germinate and within 14 days after application.
5. Is activated when the treated area receives at least 0.5 inch of irrigation or rainfall, or shallow (1 to 2 inches) mechanical incorporation.

Use Precautions

To reduce injury potential:

a. In the spring when buds are rapidly growing and expanding, over-the-top application of Cavalcade 65WDG may temporarily injure new growth of desirable plants. To reduce the possibility of injury at this time, wait to apply Cavalcade 65WDG over-the-top of newly emerged vegetation until it has hardened off, unless your experience indicates that the ornamental plant will not be injured by the other-the-top application.

b. After application, immediately irrigate the treated area to wash Cavalcade 65WDG from plant surfaces onto soil (watering plants before application may improve the washing process).

Ornamentals and Christmas Tree Farms – Application Sites and Instructions

Site	Application Instructions
Newly-Transplanted container or Field Nursery Stock	1. Delay application until soil has settled around transplants. 2. Water transplants thoroughly before application. 3. Apply after cuttings from roots are established. 4. To avoid inhibition of the tissue union, apply before budding/grafting or after buds/grafts have taken.
Established Container, Field Nursery Stock, or Landscape Plants	Apply at any time as a broadcast, over-the-top, or directed spray.
Landscape (or Ornamental) Plantings	1. Apply as a broadcast, over-the-top, or as a directed spray. 2. Delay application to newly-transplanted ornamentals until soil has settled around transplants.
Bare Ground Application for Container Placement	1. Apply to soil (including mulch, gravel, wood chips, or other permeable base) upon which containerized ornamentals are placed. 2. After Cavalcade 65WDG is applied, perform shallow cultivation or hand weeding only, or avoid disturbing the herbicide barrier.
In Shade Houses and Uncovered Polyhouses	After Cavalcade 65WDG is applied, uncovered polyhouses must remain open for at least 7 days and ornamentals must receive 2 irrigations totaling at least ½ inch of water.
Ornamental Bulbs and Perennial Wildflower Plantings	1. Cavalcade 65WDG may be applied to bulbs or perennial wildflower species listed in the section. **Tolerant Ornamental Species**. 2. Apply before or after bulbs emerge but before bulbs bloom and weeds emerge. In wildflowers, a postemergence herbicide labeled for wildflowers may be needed to control weeds that have already emerged.

How Much and When to Apply

Amount to Apply (Broadcast)*	When to Apply	Comments/Instructions
1.0-2.3 lbs./A or 0.37-0.83 oz./1,000 sq. ft.	In fall or spring before weeds germinate or after weeds are removed.	1. Use the higher rate for longer control. 2. Cavalcade 65WDG may be applied more than once per year as long as the total amount of product applied does not exceed 2.3 lbs./A per year.

*Note: For band application calculate amount per acre:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate} = \text{amount to apply per acre of field}$$

Equivalent Measurements for Cavalcade 65WDG

Lbs./A	Oz./1,000 sq. ft.	Approximate Equivalent-Tablespoons/1,000 sq. ft.
1.0	0.37	1
1.5	0.55	1.5
2.0	0.74	2
2.3	0.83	2.25

Tank Mixtures For Use On Ornamentals

Cavalcade 65WDG may be tank mixed with other registered herbicides listed on this label to provide a broader spectrum of weed control or to control emerged weeds. Tank mixes with Cavalcade 65WDG are for use only in states where the tank mix partner(s), application site, and intended use pattern are registered. Follow the label(s) of the tank mix partner(s) for application rates, timing, weeds controlled, tolerant ornamentals, and specific use precautions and/or restrictions. Before mixing pesticides in the spray tank, test compatibility by mixing the products in a small container first. See the **Compatibility Test** section of this label.

Tank Mix Partners for Cavalcade 65WDG on Ornamentals

Product	Precautions/Instructions
Goal® (use on conifers only)	Mix with Cavalcade 65WDG for postemergence control of certain broadleaf weeds including malva and filaree.
Gallery®, Sim-Trol®, Pennant®	See product labels for weed spectrum and tolerant ornamentals.
Roundup®, or other glyphosate-based products[1], Finale®	These nonselective tank mix herbicides control most emerged annual broadleaves and grasses. Take extreme care to prevent tank mixtures with these products from contacting the foliage and stems of turfgrass, trees, shrubs, or other desirable vegetation because desirable vegetation may be severely injured or killed. Apply these tank mixtures as a directed spray and use a shield to prevent spray from contacting foliage of desirable plants. Following instructions on the tank mix partner's label, delay irrigation of the treated area to allow time for the herbicide to be absorbed by weed foliage.

[1] Roundup is one brand of a nonselective herbicide containing glyphosate. Other glyphosate products may also be used.

Tolerant Ornamental Species

Cavalcade 65WDG will not harm most trees, shrubs, vines, and flowers. The species listed below in Table 1 are tolerant to Cavalcade 65WDG. Cavalcade 65WDG is approved for application, except in CA, to the species in Table 2. Cavalcade 65WDG may be applied over-the-top of the listed species.

When plants are under stress (such as heat, drought, or frost damage), some cultivars of listed plants may be sensitive to Cavalcade 65WDG.

Table 2 – Tolerant Ornamental Species – All States

Scientific name	Common name
Abelia grandiflora	Abelia
Abies spp.	Fir species (Balsam, Fraser, Noble, etc.)
Acer palmatum	Japanese maple
Acer platanoides	Norway maple
Achillea spp.	Yarrow
Actinidia chinensis	Kiwi*
Agapanthus orientalis	Lily of the Nile; African lily
Akebia quintata	Five-Leaf or Chocolate Vine

(continued)

Table 2 – Tolerant Ornamental Species – All States *(continued)*

Scientific name	Common name
Allium cernuum	Lady's Leek; Nodding Onion
Anemone hybrida	Japanese Anemone
Aquilegia spp.	Columbine
Arctostaphylos densiflora	Vine hill manzanita
Arctotheca calendula	Cape weed
Aucuba japonica	Japanese Aucuba
Artemisia spp.	Wormwood; Silver Mound; Castle
Aster spp.	Aster
Athyrium filix-femina	Lady Fern
Begonia spp.	Fibrous Begonia
Berberis gladwynesis	Barberry
Berberis julianae	Wintergreen barberry
Berberis mentorensis	Mentor barberry
Berberis thunbergii	Japanese barberry
Berberis verriculosa	Warty barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Bougainvillea spp.	Bougainvillea
Buddleia davidii	Butterfly-bush
Buxus microphylla	Japanese boxwood
Callistemon citrinus	Crimson bottlebrush
Callistemon viminalis	Weeping bottlebrush
Calluna vulgaris	Scotch heather
Campanula carpatica	Tussock bellflower
Campsis X tagliabuana	Trumpet creeper, Trumpet flower
Carpobrotus edulis	Hottentot fig; Ice plant
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild lilac
Ceratostigma plumbaginoides	
Chamaecyparis pisifera	False cypress
Chrysanthemum nipponicum	
Cleyera japonica	Cleyera
*Citrus spp.**	Ornamental orange, lemon, lime, etc.*
Coreopsis spp.	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
Cornus stolonifera	American dogwood
Cortaderia selloana	Pampas grass
Cotoneaster apiculatus	Cranberry Cotoneaster
Cotoneaster buxifolius	Cotoneaster
Cotoneaster dammeri	Bearberry Coteneaster
Cotoneaster microphyllus	Rockspray Cotoneaster
Crataegus spp.	Hawthorn
Cupressus sempervirens	Italian cypress
Crocosmia spp.	Lucifer
Delosperma spp.	Ice plant
Delphinium spp.	Larkspur
Dianthus deltoidia	Dianthus; Maiden pinks
Dianthus gratianopolitanus	Cheddar pink
Dodonea viscosa	Hop bush
Echinacea purpurea	Coneflower
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
Euonymus japonica	Japanese spindle tree; Evergreen Euonymus
Euonymus kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese aralia
Forsythia intermedia	Border Forsythia
Forsythia suspense	Weeping Forsythia
Forsythia viridissima	Greenstem Forsythia

(continued)

Table 2 – Tolerant Ornamental Species – All States *(continued)*

Scientific name	Common name
Gaillardia spp.	Gaillardia; Blanket flower
Gardenia jasminoides	Gardenia; Cape-jasmine
Gaura spp.	Gaura
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
*Gladiolus spp.***	Gladiolus species**
Gypsophila repens	Baby's breath
Hedera helix	English ivy
Helianthemum spp.	Sunrose
Hemerocallis spp.	Daylily
Heucherella spp.	Coral bells
*Hibiscus rosa-sinensis***	Chinese Hibiscus**
Hibiscus spp.	Mallow; Rose of Sharon**
Hosta plantaginea	Hosta; Plantain lily
Hosta sieboldiana	Hosta
Houttuynia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
*Ilex cornuta***	Chinese holly**
Ilex crenata	Japanese holly
Ilex opaca	American holly
Ilex pernyi	Holly
Ilex vomitoria	Yaupon holly
Inula ensifolia	
Iris spp.	Iris
Jasminium nudiflorum	Winter jasmine
*Juglans spp.**	Walnut*
Juniperus chinensis	Chinese juniper
Juniperus conferta	Shore juniper
Juniperus davurica	
Juniperus horizontalis	Creeping juniper
Justicia brandegeana	Shrimp plant
Lagerstroemia indica	Crape myrtle
Lagerstroemia indica and hybrids	Crape myrtle
Lantana montevidensis	Weeping Lantana
Lavender spp.	Lavender; Munstead
Leontopodium alpinum	Edelweiss
Ligustrum amurense	Amur privet
Ligustrum japonicum	Japanese privet
Ligustrum lucidum	Glossy privet; Wax-leaf
Ligustrum sinense	Chinese privet
Lilium spp.	Lily
Liriope muscari	Liriope
Liriope spicata	Liriope, creeping
Lobelia cardinalis	Cardinal flower; Indian pink
Lonicera japonica	Japanese honeysuckle
Lonicera tatarica	Tatarian honeysuckle
Loropetalum chinense	Loropetalum
Lythrum spp.	Loosestrife
*Magnolia spp.***	Magnolia
Maleophora luteola	Ice plant
*Malus spp.**	Crabapple*
*Miscanthus sinesis***	Yaku Jima**, Silberfeder**
Nandina domestica	Heavenly bamboo
*Narcissus spp.***	Narcissus, Daffodil
Nerium spp.	Oleander
Oenothera missouriensis	Evening primrose

(continued)

Table 2 – Tolerant Ornamental Species – All States *(continued)*

Scientific name	Common name
*Olea europaea**	Olive*
*Ophiopogon japonicus***	Mondo grass**
Osmanthus heterophyllus	Osmanthus; False holly
Osteospermum fruticosum	Trailing African daisy
Oxydendron luteum	Sourwood
Paeonia suffruticosa	Tree peony
*Pennisetum setaceum***	Fountain grass**
Perovskia atriplicifolia	
*Persea americana**	Avocado*
Photinia fraseri	Photinia; Redtip
Physostegia virginiana	False dragonhead
*Picea spp.***	Spruces (Colorado Blue, Norway, etc.)
Pieris japonica	Japanese andromeda; Lily-of-the-valley shrub
Pinus brutia	Calabrian pine
Pinus canariensis	Canary island pine
Pinus elliottii	Slash pine
Pinus halepensis	Aleppo pine
Pinus nigra	Austrian black pine
Pinus palustris	Longleaf pine
Pinus radiata	Monterey pine
Pinus strobus	Eastern white pine
Pinus sylvestris	Scotch pine
Pinus taeda	Loblolly pine
Pinus thunbergiana	Japanese black pine
Pinus virginiana	Virginia pine
*Pistacia spp.**	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum
Pittosporum tobira	Japanese Pittosporum
Podocarpus macrophyllus	Japanese yew
Prunus laurocerasus	English laurel
*Prunus spp.**	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
*Pseudotsuga menziesii***	Douglas fir**
Pyracantha coccinea	Firethorn, scarlet
Pyracantha fortuneana	Firethorn
Pyracantha koidzumii	Firethorn
Pyrus spp.	Pear spp., including 'Bradford'
Quercus rubra	Red oak
Quercus shumardii	Shumard oak
Raphiolepsis indica	Indian hawthorne
Raphiolepsis umbellata	Yedda hawthorne
Rhododendron spp.	Rhododendrons, Azaleas
Rosa banksiae	Lady Banks rose
Rudbeckia spp.	Black-eyed Susan
Rumohra adiantiformis	Leatherleaf Fern
Santolina virens	
Saxifraga spp.	Saxifrage; Purple dome
Scabiosa spp.	Pincushion flower
Sedum spp.	Stonecrop
Spiraea bumalda	Spirea
Syzygium paniculatum	Australian brushcherry; Japanese boxcherry
Taxus cuspidata	Japanese yew
Taxus spp.	Yew
Teucrium spp.	Germander
Thalictrum dipterocarpum	Meadow rue
Thuja occidentalis	American arborvitae
Trachelospermum asiatum	Star jasmine
Tsuga canadensis	Canada hemlock

(continued)

Table 2 – Tolerant Ornamental Species – All States *(continued)*

Scientific name	Common name
Tulipa spp.	Tulip
Veronica spp.	Veronica; Speedwell
Viburnum japonicum	Japanese viburnum
Viburnum odoratissimum	Sweet viburnum
Viburnum plicatum	Japanese snowball
Viburnum rigidum	Canary island viburnum
Viburnum japonicum	Japanese viburnum
Viburnum suspensum	Arrowood viburnum
Viburnum tinus	Laurustinus
Viburnum trilobium	Cranberry bush
Viburnum wrightii	Leatherleaf viburnum
Vinca major	Vinca
Vinca minor	Periwinkle
*Vitis spp.**	Grape*
Weigela florida	Old fashioned Weigela
Yucca aloifolia	Spanish bayonet
Yucca filamentosa	Yucca; Adam's needle

* Do not use on food producing trees, vines, or plants.
** Not for use on container grown plants.

NEW PLANTINGS, REPLANTING, AND ROTATIONAL PLANTINGS

Nursery, landscape, or non-cropland areas treated with Cavalcade 65WDG should be rotated only to ornamental species listed on this label for 1 year following application unless the following test has shown species safety:

Before planting a species not listed on this label, it is recommended that several test strips of an indicator plant such as wheat, sorghum, or corn be sown into the treated area. If the indicator plants germinate and grow normally to a height of 12 inches with normal root development, it is safe to plant.

In areas disturbed by new plantings or replanting of labeled species, it may be necessary to retreat exposed soil to maintain satisfactory weed control.

CHEMIGATION INSTRUCTIONS – OVERHEAD SPRINKLER IRRIGATLON APPLICATION

- Apply this product only through an overhead sprinkler irrigation system. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
- To avoid injury to foliage, make sure foliage is sufficiently wet before application or adequate irrigation is applied after application.
- If sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result.
- If sprinkler distribution patterns overlap excessively, injury to leatherleaf ferns and other ornamentals may result.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to public water systems unless pesticide label-prescribed safety devices for public water systems are in place.
- If necessary, a person knowledgeable of the chemigation system and responsible for its operation, or someone under the supervision of the responsible person, shall shut the system down arid make necessary adjustments.

Operation Instructions

1. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water. pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump such as a positive displacement injection pump (e g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Prepare a mixture with a minimum of 20 parts of water to 1 part Cavalcade 65WDG and inject this herbicide suspension mixture into the overhead system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
9. Before injecting Cavalcade 65WDG in to the system, run the irrigation system long enough to wet the foliage, then inject Cavalcade 65WDG suspension mixture in the pesticide supply tank (see number 8 above) in 1 inch of irrigation water. After the application is complete, continue the irrigation until all residues are washed off the foliage.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store in original container away from feed or foodstuffs and separated from other pesticides.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

CONTAINER DISPOSAL:

Paper and plastic film bags or boxes- Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or incinerate; or, if allowed by state and local authorities, burn locally. Stay out of smoke from burning container.

Fiber drums with plastic liners- Completely empty the plastic liner by shaking and tapping sides and bottom to loosen clinging particles. Pour residues into application equipment. Dispose of empty liner at an incineration facility. Offer the fiber drum for recycling or re-use. If the fiber drum cannot be recycled or re-used, dispose of at an incineration facility.

Plastic containers- Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call CHEMTREC (phone: 1-800-424-9300) day or night.

WARRANTY AND LIMITATION OF DAMAGES

CONDITIONS OF SALE: Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc.. **SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.**

Sim-Trol is a trademark of Sipcam Agro USA, Inc.
Pennant is a trademark of Syngenta Crop Protection Co.
Finale is a trademark of Bayer CropScience
Gallery and Goal are trademarks of Dow AgroSciences
Roundup is a trademark of Monsanto Company

SIPCAM AGRO USA, INC.

CAVALCADE™ 65WDG

For preemergence control of grass and broadleaf weeds in:

- Established turfgrasses (excluding golf course putting greens), lawns and sod nurseries
- Landscape ornamentals in nurseries or in established plantings
- Established perennials and wildflower plantings
- Plants grown for cut foliage production (Florida only)
- Conifer and hardwood tree seedling nurseries
- Christmas tree farms
- Managed transportation and utility rights-of-way, including rail and equipment yards, and public utility facilities (substations, tank farms, pumping stations, parking/storage areas, ungrazed fencerows)

ACTIVE INGREDIENT:	
Prodiamine (CAS No. 29091-21-2)	65.0%
OTHER INGREDIENTS:	35.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have affected person sip a glass of water if able to swallow. • Do not induce vomiting unless told by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 900-4044 Poison Control Center (human health) (800) 345-4735 ASPCA (animal health)

Cavalcade is a trademark of Sipcam Agro USA, Inc.

SIPCAM AGRO USA, INC.
300 COLONIAL CENTER PARKWAY, SUITE 230
ROSWELL, GA 30076

EPA REG. NO. 60063-26
EPA EST. NO. 37429-GA-01

NET CONTENTS: 10 pounds