

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Non-refillable container. Do not reuse or refill container.

Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

STEP 2



STEP 1



MADE IN THE USA

EPA Reg. No.: 3282-65
EPA Est. No.: 475-MS-1

SATISFACTION GUARANTEED
OR YOUR MONEY BACK

THIS SIDE UP. PULL TAB.
LIFT AND TWIST OFF.



HOUSE MICE MAY CONSUME
A LETHAL DOSE IN ONE
FEEDING WITH FIRST DEAD
MICE APPEARING 4 OR 5 DAYS
AFTER FEEDING BEGINS.

KEEP OUT OF REACH
OF CHILDREN

CAUTION:

May be harmful or fatal if swallowed.
Read additional precautionary statements on outer panel.

NET CONTENTS: 1.5 OZ. (43g)

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-1,1'-biphenyl)-4-yl]-
-1,2,3,4-tetrahydro-1-naphthalenyl] 0.005%
4-hydroxy-2H-1-benzopyran-2-one 0.005%

OTHER INGREDIENTS: 99.995%

TOTAL: 100.000%

1

Bait Wedge / Triángulo con cebo



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. **9833.114**
Disc.

IMPORTANTE: Para instrucciones de uso y primeros auxilios en español por favor llame al 1-866-648-1819.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

USE RESTRICTIONS: THIS IS NOT A TAMPER-PROOF OR TAMPER RESISTANT BAIT STATION. DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS, OR NONTARGET WILDLIFE.

DO NOT USE THIS PRODUCT OUTDOORS.

This product may be used to control house mice in homes. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Wedges of d-CON® Mouse-Prufe II may be placed indoors along walls, in corners and in concealed places where mice are active and where children, nontarget mammals and birds will not be exposed to the bait.

APPLICATION DIRECTIONS: Place wedge in horizontal position. Pull tab, lift and twist off. Space wedges at intervals of 8 to 12 feet if mouse activity is not highly localized and placements can be made in areas not accessible to children and nontarget animals. Replace wedges as their contents are consumed. Wedges not touched after 3 days should be moved to new locations appropriate for bait placements (see **USE RESTRICTIONS**); if mouse activity persists. Maintain a supply of fresh bait for at least 15 days. Collect and dispose of all dead animals and leftover bait properly.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Wash hands after handling bait.

FIRST AID

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE.

IF SWALLOWED BY HUMAN

- Call a Poison Control Center, doctor or 1-800-228-4722 immediately for treatment advice.
- Do not induce vomiting unless told to do so by the Poison Control Center or doctor.

IF SWALLOWED BY PET OR ANIMAL

- Call a veterinarian immediately for treatment advice.

NOTE TO PHYSICIAN AND VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.



Beckitt
Beckitt

Questions? Comments?
Call 1-800-228-4722
www.beckitt.com
Distributed by: Beckitt Beckitt
Parapary, NJ 07054-0224
Made in USA © RB 2011

0342402

2

19200 00051

0

THIS SIDE UP. PULL TAB.
LIFT AND TWIST OFF.

d-CON[®]

Mouse-Prufe II

KILLS MICE

MATA RATONES

**HOUSE MICE MAY CONSUME
A LETHAL DOSE IN ONE
FEEDING WITH FIRST DEAD
MICE APPEARING 4 OR 5 DAYS
AFTER FEEDING BEGINS.**

KEEP OUT OF
REACH OF CHILDREN

CAUTION:

May be harmful or fatal if swallowed
Read additional precautionary statements on back panel.

NET CONTENTS: 3.0 OZ. (85 g)

1

Bait Wedge/Triángulo con cebo

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-1,1'-biphenyl-4-yl)
-1,2,3,4-tetrahydro-1-naphthalenyl]
-4-hydroxy-2H-1-benzopyran-2-one 0.005%

OTHER INGREDIENTS: 99.995%

TOTAL: 100.000%



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

IMPORTANTE: Para instrucciones de uso y primeros auxilios en español por favor llame al 1-866-648-1819.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

USE RESTRICTIONS: THIS IS NOT A TAMPER-PROOF OR TAMPER RESISTANT BAIT STATION. DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS, OR NONTARGET WILDLIFE. DO NOT USE THIS PRODUCT OUTDOORS.

This product may be used to control house mice in homes. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Wedges of d-CON® Mouse-Prufe II may be placed indoors along walls, in corners and in concealed places where house mice are active and where children, nontarget mammals and birds will not be exposed to the bait.

APPLICATION DIRECTIONS: Place wedge in horizontal position. Pull tab, lift and twist off.

Space wedges at intervals of 8 to 12 feet if mouse activity is not highly localized and placements can be made in areas not accessible to children and nontarget animals. Replace wedges as their contents are consumed.

Wedges not touched after 3 days should be moved to new locations appropriate for bait placements (see **USE RESTRICTIONS**.) if mouse activity persists. Maintain a supply of fresh bait for at least 15 days. Collect and dispose of all dead animals and leftover bait properly.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Wash hands after handling bait.

**FIRST AID
HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE.**

IF SWALLOWED BY HUMAN	<ul style="list-style-type: none">• Call a Poison Control Center, doctor or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the Poison Control Center or doctor.
IF SWALLOWED BY PET OR ANIMAL	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.

NOTE TO PHYSICIAN AND VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bistydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Non-refillable container. Do not reuse or refill container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.




MADE IN THE USA

SATISFACTION GUARANTEED OR YOUR MONEY BACK



0342407

Questions? Comments? Call 1-800-228-4722
www.dbaiafo.com
Distributed by: Reckitt Benckiser
Parsippany, NJ 07054-0224
Made in USA © RB 2011



d-CON

Mouse-Prufe II

KILLS

MICE

MATA RATONES




d-CON

Mouse-Prufe II

KILLS

MICE

MATA RATONES



STEP 1



STEP 2

**SATISFACTION GUARANTEED
OR YOUR MONEY BACK**

**HOUSE MICE MAY CONSUME
A LETHAL DOSE IN ONE
FEEDING WITH FIRST DEAD
MICE APPEARING 4 OR 5 DAYS
AFTER FEEDING BEGINS.**

**KEEP OUT OF
REACH OF CHILDREN
CAUTION:**

May be harmful or fatal if swallowed.
Read additional precautionary
statements on back panel.

**NET CONTENTS:
4/1.5 OZ.
NET WT.: 6 OZ. (170g)**



ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-1,1'-biphenyl)-4-yl]	
-1,2,3,4-tetrahydro-1-naphthalenyl]	
-4-hydroxy-2H-1-benzopyran-2-one	0.005%
OTHER INGREDIENTS:	99.995%
TOTAL:	100.000%

d-CON

Mouse-Prufe II

KILLS

MICE

MATA RATONES



NOTICE TO BUYER AND USER:

Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.



MADE IN THE USA

**EPA Reg. No.: 3282-65
EPA Est. No.: 475-MS-1**

4 Bait Wedges
Triángulos con cebo

4 Bait Wedges
Triángulos con cebo

4 Bait Wedges
Triángulos con cebo

Questions? Comments?
Call 1-800-228-4722
www.rbna.info.com
Distributed by:
Reckitt Benckiser
Parsippany, NJ 07054-0224
Made in USA © RB 2011



0342405

INSERT
90%
UPC

0 19200 00948 5



STATE OF HAWAII
Department of Agriculture

ACCEPTED
LICENSE NO. **9833.114**
Disc.

IMPORTANTE:

Para instrucciones de uso y primeros auxilios en español por favor llame al 1-866-648-1819.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

USE RESTRICTIONS: **THIS IS NOT A TAMPER-PROOF OR TAMPER RESISTANT BAIT STATION. DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS OR NONTARGET WILDLIFE. DO NOT USE THIS PRODUCT OUTDOORS.**

This product may be used to control house mice in homes. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Wedges of d-CON® Mouse-Prufe II may be placed indoors along walls, in corners and in concealed places where mice are active and where children, nontarget mammals and birds will not be exposed to the bait.

APPLICATION DIRECTIONS: Place wedge in horizontal position. Pull tab, lift and twist off. Space wedges at intervals of 8 to 12 feet if mouse activity is not highly localized and placements can be made in areas not accessible to children and nontarget animals. Replace wedges as their contents are consumed. Wedges not touched after 3 days should be moved to new locations appropriate for bait placements (see **USE RESTRICTIONS:**) if mouse activity persists. Maintain a supply of fresh bait for at least 15 days. Collect and dispose of all dead animals and leftover bait properly.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets.

Wash hands after handling bait.

FIRST AID

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE.

IF SWALLOWED
BY HUMAN

- Call a Poison Control Center, doctor or 1-800-228-4722 immediately for treatment advice.
- Do not induce vomiting unless told to do so by the Poison Control Center or doctor.

IF SWALLOWED BY
PET OR ANIMAL

- Call a veterinarian immediately for treatment advice.

NOTE TO PHYSICIAN AND VETERINARIAN

The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin).

FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: **If empty:** Non-refillable container. Do not reuse or refill container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.