1.5"

Add color targets/marks here

20mm zipper

DISCONTINUED

d-CON®

Bait Pellets II

KILLS MICE

MATA RATONES

STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO.
9833.115

HOUSE MICE MAY CONSUME A LETHAL DOSE IN ONE FEEDING WITH FIRST DEAD MICE APPEARING 4 OR 5 DAYS AFTER FEEDING BEGINS.

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be harmful or fatal if swallowed.
Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:
Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)
-1,2,3,4-tetrahydro-1-naphthalenyl]
-4-hydroxy-2H-1-benzopyran-2-one.....0.005%
OTHER INGREDIENTS 99.995%
TOTAL: 100.000%

NET CONTENTS: 64/1.0 OZ. (28g)
Bait Packs
TOTAL: 64 OZ. (4LB.) 1.8 kg

64 Place Packs
Sobres con cebo

.375" Seal Area - No Critical Art

Trim Area - Fill with background color

3.75"

20mm zipper

STEP 1

STEP 2

House mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

IMPORTANTE: Para instrucciones de uso y primeros auxilios en español por favor llame al 1-866-648-1819.

KILLS WARFARIN-RESISTANT HOUSE MICE

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL:

Read this entire label and follow all use directions and use precautions.

IMPORTANT:

Do not expose children, pets, or other non-target animals to rodenticides.

To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.

2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.

3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice in and around homes. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where mice will most likely find and consume the bait. Generally,these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors or walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:")

APPLICATION DIRECTIONS:

TO CONTROL HOUSE MICE: Use one bait pack per placement.

Space placements at intervals of 8 to 12 feet in infested areas.
Up to 2 bait packs may be used at points of very high mouse activity.
Maintain an uninterrupted supply of fresh bait for at least 15 days.
Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by mice for 5 consecutive days, they may be relocated to other places where mice activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait station and replenish as needed.

FIRST AID HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE.	
IF SWALLOWED BY HUMAN	• Call a Poison Control Center, doctor or 1-800-228-4722 immediately for treatment advice. • Do not induce vomiting unless told to do so by the Poison Control Center or doctor.
IF SWALLOWED BY PET OR ANIMAL	• Call a veterinarian immediately for treatment advice
NOTE TO PHYSICIANS AND VETERINARIAN	
This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K_1 are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K_1 is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K_1 is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K_1 should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.	

PRECAUTIONARY STATEMENTS:
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Non-refillable container. Do not reuse or refill container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

.375" Seal Area - No Critical Art

Trim Area - Fill with background color

.375" Seal Area - No Critical Art

Trim Area - Fill with background color

NOTICE TO BUYER AND USER:
Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

LIMITED WARRANTY TO CONSUMERS
Good Housekeeping Promises
REPLACEMENT OR REFUND IF DEFECTIVE

SATISFACTION GUARANTEED
OR YOUR MONEY BACK

Questions? Comments?
Call 1-800-228-4722
www.rbnainfo.com
Distributed by: Reckitt Benckiser
Parsippany, NJ 07054-0224
Made in USA © RB 2011
EPA Reg. No.: 3282-74,
EPA Est. No.: 475-MS-1

Reckitt Benckiser

INSERT
90%
UPC
0 19200 84048 4

038564 6

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. **IMPORTANT:** Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents: 1. Store product not in use in a location out of reach of children and pets. 2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooded livestock, raccoons, bears, other potentially destructive animals or in areas prone to vandalism. 3. Dispose of product container, and unused, spoiled and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice in and around homes. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen.

Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

APPLICATION DIRECTIONS: To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days. Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by mice for 5 consecutive days, they may be relocated to other places where mice activity exists and where bait placements consistent with the requirements of this label can be made. Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

ENVIRONMENTAL HAZARDS:
This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals.
Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.
DISPOSAL: If empty: Non-refillable container. Do not reuse or refill container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Questions? Comments? Call 1-800-228-4722
www.dconinfo.com
Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in USA © RB 2011 0342408

d-CON® Bait Pellets II KILLS MICE

House Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins

Good Housekeeping Promises — LIMITED WARRANTY TO CONSUMERS — REPLACEMENT OR REFUND IF DEFECTIVE

Bait Pellets II

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be harmful or fatal if swallowed. Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:
Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one..... 0.005%
OTHER INGREDIENTS: 99.995%
TOTAL: 100.000%

EPA Reg.No. 3282-74
EPA Est.No. 475-MS-1

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

PRECAUTIONARY STATEMENTS:
HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals,and pets. Wash hands after handling bait.

First Aid
Have Label With You When Obtaining Treatment Advice.

IF SWALLOWED BY HUMAN	• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice. • Do not induce vomiting unless told to do so by the Poison Control Center or doctor.
IF SWALLOWED BY PET OR ANIMAL	• Call a veterinarian immediately for treatment advice.

NOTE TO PHYSICIAN AND VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K_1 are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K_1 is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K_1 is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K_1 should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

NET CONTENTS: 1.0 OZ (28 g)

Reckitt Benckiser

INSERT 80% UPC 19200 00009 3