DO NOT TEAR OPEN THE INDIVIDUAL PACK - THE MICE WILL CHEW THROUGH THE WRAPPER TO GET AT THE BAIT.

HOUSE MICE CAN CONSUME A LETHAL DOSE IN ONE NIGHT'S FEEDING WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL. and follow all use directions and precautions.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be taken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooted livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may only be used to control house mice. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment. Do not broadcast bait.

Urban areas: This product may be used in and around homes, food processing facilities, industrial, commercial, agricultural and public buildings, and similar man-made structures. It may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. This product may also be used in alleys.

Non-urban areas: This product may only be used inside of homes and agricultural, industrial and commercial buildings.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume bait. Generally, these are along walls, by gnawed openings, in corners and concealed places, between floors or walls or in locations where rodents or their signs have been observed. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Mice: Apply 1 pack per placement, usually spaced 8 to 12 feet apart. Two packs may be needed at points of very high mouse activity. Maintain a constant supply of fresh bait for 15 days or until signs of mouse activity cease.

Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed.

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

CAUTION: Keep away from children, domestic animals and pets. May be harmful if swallowed or absorbed through the skin because it may reduce the clotting ability of blood and cause bleeding. Avoid contact with eyes, skin or clothing. Wear gloves when handling bait or animals. Wash arms, hands and face with soap and water after applying and before eating and smoking.

FIRST AID

Have this label with you when obtaining treatment advice.

If swallowed	• Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin with plenty of cool water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.

NOTE TO PHYSICIAN OR VETERINARIAN

Contains difethialone, an anticoagulant. For humans or dogs that have ingested this product, or have obvious poisoning symptoms (bleeding) or prolonged prothrombin times, give Vitamin K_1 by intramuscular or oral administration. Check prothrombin time every 3 days until values return to normal.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

STORAGE AND DISPOSAL: This is a nonrefillable container.

Storage: Store in original container in a dry place inaccessible to children and pets.

Disposal: If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

INSERT
100%
UPC
0 19200 80878 1

EPA Reg. No. 7173-258-3282
EPA Est. No. 7173-WI-1
0166560 100-3851-0408

Questions?
Call: 1-800-228-4722

Distributed by:
Reckitt Benckiser Inc.,
Parsippany, NJ 07054-0224

©RBI 2008 Made in France

SATISFACTION GUARANTEED OR YOUR MONEY BACK

9833.210

DISC.

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD **2008-2010** LIC. NO.

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD **2014-2016** LIC. NO.

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD **2011-2013** LIC. NO.

d-CON®

Bait Paste Pouches™

KILLS HOUSE MICE

MATA RATONES CASEROS

DEMONSTRATED PALATABILITY FOR MICE

KEEP OUT OF REACH OF CHILDREN
CAUTION: See back panel for additional precautionary statements.

NET WT.: 2.5 OZ. (7x10g)

ACTIVE INGREDIENT:
difethialone..... 0.0025%
INERT INGREDIENTS:..... 99.9975%
TOTAL:..... 100.0000%

7 Bait Paste Pouches™