

- Germicidal • Virucidal* • Fungicidal • Tuberculocidal • Bactericidal • Mildewcidal • Kills antibiotic resistant bacteria*
- Kills the source of mold allergens and prevents it from coming back for up to 7 days††
- Deodorizes by killing odor-causing bacteria at their source. Ordinary, non-germicidal sprays can't do this.
- Use on soft surfaces where bacteria can hide*** Meets OSHA Bloodborne Pathogens Standards§§
- Meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants
- ***Kills 99.9% of Staphylococcus aureus and Klebsiella pneumoniae on soft surfaces in 30 seconds

| | | | |
|--|--|--|--|
| Kills in 30 seconds: | | | |
| • Avian Influenza A Virus | • Hepatitis C Virus | • Type 1 [HIV-1] [AIDS Virus] | • Rhinovirus Type 39† |
| • Campylobacter jejuni | • Herpes Simplex Virus Type 1 | • Influenza A Virus* | • Rotavirus WA** |
| • Coxsackie Type B3 Virus | • Herpes Simplex Virus Type 2 | • Influenza B Virus | • SARS Coronavirus |
| • Cytomegalovirus | • Human Coronavirus | • Newcastle Disease Virus | • Swine Influenza A Virus |
| • Hepatitis B Virus | • Human Immunodeficiency Virus | • Respiratory Syncytial Virus (RSV)** | • Vaccinia Virus |
| Kills in 2 minutes: | | | |
| • Acinetobacter baumannii MDR | • Enterovirus Type 68 | • Proteus mirabilis | • Staphylococcus haemolyticus |
| • Bordetella pertussis | • Escherichia coli | • Proteus vulgaris | • Staphylococcus saprophyticus |
| • Burkholderia cepacia | • Haemophilus influenzae | • Pseudomonas aeruginosa** | • Streptococcus pneumoniae PRSP |
| • Candida albicans | • Klebsiella pneumoniae | • Salmonella enterica** | • Streptococcus pyogenes (Strep) |
| • Candida parapsilosis | • Klebsiella pneumoniae NDM-1 positive | • Serratia marcescens | • Streptococcus salivarius (Pink Eye) |
| • Citrobacter freundii | • Klebsiella pneumoniae Carbapenem resistant | • Shigella dysenteriae | • Trichophyton mentagrophytes (Athlete's Foot Fungus) (Ringworm) |
| • Corynebacterium diphtheriae | • Listeria monocytogenes | • Shigella flexneri | • Vibrio cholerae |
| • Cronobacter sakazakii | • Neisseria elongata | • Shigella sonnei | • Yersinia enterocolitica |
| • Enterobacter aerogenes MDR | • Pasteurella multocida | • Staphylococcus aureus** | |
| • Enterococcus faecalis | • Poliovirus Type 1 | • Staphylococcus aureus - Methicillin resistant (HA-MRSA)* | |
| • Enterococcus faecium VRE + MDR | | • Staphylococcus epidermidis | |
| Kills in 3 minutes: • Adenovirus 14 • Aspergillus niger† | | | |
| Kills in 10 minutes: • Feline Calicivirus (Norovirus) • Mycobacterium bovis BCG (Quant tuberculosis) • Bordetella avium | | | |

Intended for use in:

- Ambulances • Clinics • Dental Offices • Hotel/Motel • Office Buildings • Physicians Offices • Schools
- Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices
- Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

| | | | | | | |
|----------------------------------|----------------------|--------------------|------------------|--------------------|----------------------|--------------|
| Hard, non-porous surfaces | • Bathroom Surfaces | • Door Handles | • Light Switches | • Sports Equipment | Soft surfaces | • Bedding |
| • Chairs | • Examination Tables | • Portable Toilets | • Shower Areas | • Tables | | • Couches |
| • Counters | • Floors | • Telephone | • Toilet Seats | • Curtains | | • Mattresses |
| • Desks | • Garbage Cans | • Sinks | | | | |
| | | | | | | |

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

To Disinfect: Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes then allow to air dry. **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry. **Fabric Sanitizer:** To Sanitize Soft Surfaces: Spray until fabric is wet. DO NOT SATURATE. Fabric must remain wet for 30 seconds. Let air dry. For difficult odors, repeat application.

Rinse toys and food contact surfaces with potable water after use.

To Control and Prevent Mold & Mildew: Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears. **To Deodorize:** Spray on surfaces as needed. Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

For Tuberculocidal Activity: This product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.

***KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1 AIDS VIRUS), HEPATITIS B VIRUS (HBV), AND HEPATITIS C VIRUS (HCV) ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids; and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS), Hepatitis B Virus, and Hepatitis C Virus.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV, AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS: PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1, HBV, and HCV. For other organisms, surfaces must remain wet for 3 minutes. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

CONTACT NUMBER: Questions? Comments or In case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

PHYSICAL HAZARDS: FLAMMABLE: Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed. **DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. This can be made from an average of 25% recycled steel (10% post-consumer).

Questions? Call 1-800-228-4722

For ingredient and other information, visit www.rmainfo.com

Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in U.S.A. © 2016 RB

EPA Reg. No.: 777-99-675

EPA Est. No.: 777-741-2 (See Bottom for Lot/Date code)

8232787




PLEASE RECYCLE WHEN EMPTY



FOR COMMERCIAL USE

KILLS COLD & FLU VIRUSES†



KILLS 99.9% OF VIRUSES & BACTERIA**

Disinfectant Spray

Eliminates odors

Country scent®



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. 9833.249

NET WT.:
19 OZ.
(1 LB. 3 OZ.)
538 g

ACTIVE INGREDIENTS:

Alkyl (50% C14, 40% C12, 10% C16)
dimethyl/benzyl ammonium saccharinate.....0.10%
Ethanol.....58.00%
OTHER INGREDIENTS.....41.90%
TOTAL.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS.

- Germicidal • Virucidal* • Fungicidal • Tuberculocidal • Bactericidal • Mildewcidal • Kills antibiotic resistant bacteria*
- Kills the source of mold allergens and prevents it from coming back for up to 7 days**
- Deodorizes by killing odor-causing bacteria at their source. Ordinary, non-germicidal sprays can't do this.
- Use on soft surfaces where bacteria can hide*** Meets OSHA Bloodborne Pathogens Standards³³
- Meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants
- ***Kills 99.9% of Staphylococcus aureus and Klebsiella pneumoniae on soft surfaces in 30 seconds

| | | | |
|-----------------------------|--|--|--|
| Kills in 30 seconds: | • Hepatitis C Virus • Herpes Simplex Virus Type 1 • Herpes Simplex Virus Type 2 • Human Coronavirus • Human Immunodeficiency Virus | Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus† • Influenza B Virus • Newcastle Disease Virus • Respiratory Syncytial Virus (RSV)** | • Rhinovirus Type 39† • Rotavirus WA** • SARS Coronavirus • Swine Influenza A Virus • Vaccinia Virus |
| Kills in 2 minutes: | • Acinetobacter baumannii MDR • Bordetella pertussis • Burkholderia cepacia • Candida albicans • Candida parapsilosis • Citrobacter freundii • Corynebacterium diphtheriae • Cronobacter sakazakii • Enterobacter aerogenes MDR • Enterococcus faecalis • Enterococcus faecium VRE + MDR | • Enterovirus Type 68 • Escherichia coli • Haemophilus influenzae • Klebsiella pneumoniae • Klebsiella pneumoniae NDM-1 positive • Klebsiella pneumoniae Carbapenem resistant • Listeria monocytogenes • Neisseria elongata • Pasteurella multocida • Poliovirus Type 1 | • Proteus mirabilis • Proteus vulgaris • Pseudomonas aeruginosa** • Salmonella enterica** • Serratia marcescens • Shigella dysenteriae • Shigella flexneri • Shigella sonnei • Staphylococcus aureus** • Staphylococcus aureus - Methicillin resistant (HA-MRSA)* • Staphylococcus epidermidis |
| Kills in 3 minutes: | • Adenovirus 14 | • Aspergillus niger†† | • Bordetella avium |
| Kills in 10 minutes: | • Feline Calicivirus (Norovirus) | • Mycobacterium bovis BCG (Quant tuberculosis) | |

Intended for use in:

- Ambulances • Clinics • Dental Offices • Hotel/Motel • Office Buildings • Physicians Offices • Schools
- Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices
- Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

| | | | | | | |
|----------------------------------|--|--|---|--|----------------------|--|
| Hard, non-porous surfaces | • Bathroom Surfaces • Chairs • Counters • Desks | • Door Handles • Examination Tables • Floors • Garbage Cans | • Light Switches • Portable Toilets • Shower Areas • Sinks | • Sports Equipment • Tables • Telephones • Toilet Seats | Soft surfaces | • Bedding • Couches • Curtains • Mattresses |
|----------------------------------|--|--|---|--|----------------------|--|

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

To Disinfect: Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes then allow to air dry. **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry. **Fabric Sanitizer:** To Sanitize Soft Surfaces: Spray until fabric is wet. DO NOT SATURATE. Fabric must remain wet for 30 seconds. Let air dry. For difficult odors, repeat application.

Rinse toys and food contact surfaces with potable water after use.

To Control and Prevent Mold & Mildew: Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals when mold and mildew growth appears. **To Deodorize:** Spray on surfaces as needed. Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

For Tuberculocidal Activity: This product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.

****KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1 AIDS VIRUS), HEPATITIS B VIRUS (HBV), AND HEPATITIS C VIRUS (HCV) ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids; and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS), Hepatitis B Virus, and Hepatitis C Virus.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV, AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD / BODY FLUIDS: PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1, HBV, and HCV. For other organisms, surfaces must remain wet for 3 minutes. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

CONTACT NUMBER: Questions? Comments or in case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

PHYSICAL HAZARDS: FLAMMABLE: Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed. **DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty; Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. This can is made from an average of 25% recycled steel (10% post-consumer).

Questions? Call 1-800-228-4722

For ingredient and other information, visit www.rhainfo.com

Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in U.S.A. © 2016 RB

EPA Reg. No.: 777-99-675

EPA Est. No.: 777-AU-2 (See Bottom for Lot/Date code)



8232788



PLEASE RECYCLE WHEN EMPTY




9

0 36241 74828

FOR COMMERCIAL USE

KILLS COLD & FLU VIRUSES†



KILLS 99.9% OF VIRUSES & BACTERIA**

Disinfectant Spray

Eliminates odors

Crisp Linen®
scent



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. **9833.249**

NET WT.:
19 OZ.
(1 LB. 3 OZ.)
538 g

ACTIVE INGREDIENTS:

Alkyl (50% C14, 40% C12, 10% C16)
dimethyl benzyl ammonium saccharinate.....0.10%
Ethanol.....58.00%
OTHER INGREDIENTS:.....41.90%
TOTAL:.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

- Germicidal • Virucidal† • Fungicidal • Tuberculocidal • Bactericidal • Mildewcidal • Kills antibiotic resistant bacteria*
- Kills the source of mold allergens and prevents it from coming back for up to 7 days††
- Deodorizes by killing odor-causing bacteria at their source. Ordinary, non-germicidal sprays can't do this.
- Use on soft surfaces where bacteria can hide*** Meets OSHA Bloodborne Pathogens Standards§§
- Meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants
- *** Kills 99.9% of Staphylococcus aureus and Klebsiella pneumoniae on soft surfaces in 30 seconds

| | | | |
|-----------------------------|---|--|---|
| Kills in 30 seconds: | • Hepatitis C Virus • Herpes Simplex Virus Type 1 • Herpes Simplex Virus Type 2 • Human Coronavirus • Human Immunodeficiency Virus | Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus† • Influenza B Virus • Newcastle Disease Virus • Respiratory Syncytial Virus (RSV)** | • Rhinovirus Type 39† • Rotavirus WA** • SARS Coronavirus • Swine Influenza A Virus • Vaccinia Virus |
| Kills in 2 minutes: | • Enterovirus Type 68 • Acinetobacter baumannii MDR • Bordetella pertussis • Burkholderia cepacia • Candida albicans • Candida parapsilosis • Citrobacter freundii • Corynebacterium diphtheriae • Cronobacter sakazakii • Enterobacter aerogenes MDR • Enterococcus faecalis • Enterococcus faecium VRE + MDR | • Proteus mirabilis • Proteus vulgaris • Pseudomonas aeruginosa** • Salmonella enterica** • Serratia marcescens • Shigella dysenteriae • Shigella flexneri • Shigella sonnei • Staphylococcus aureus** • Staphylococcus aureus - Methicillin resistant (HA-MRSA)* • Staphylococcus epidermidis | • Staphylococcus haemolyticus • Staphylococcus saprophyticus • Streptococcus pneumoniae PRSP • Streptococcus pyogenes (Strep) • Streptococcus salivarius (Pink Eye) • Trichophyton mentagrophytes (Athlete's Foot Fungus) (Ringworm) • Vibrio cholerae • Yersinia enterocolitica |
| Kills in 3 minutes: | • Adenovirus 14 | • Aspergillus niger† | • Bordetella avium |
| Kills in 10 minutes: | • Feline Calicivirus (Norovirus) | • Mycobacterium bovis BCG (Quant tuberculosis) | |

Intended for use in:

- Ambulances • Clinics • Dental Offices • Hotel/Motel • Office Buildings • Physicians Offices • Schools
- Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices
- Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

| | | | | | | |
|----------------------------------|--|--|---|--|----------------------|--|
| Hard, non-porous surfaces | • Bathroom Surfaces • Chairs • Counters • Desks | • Door Handles • Examination Tables • Floors • Garbage Cans | • Light Switches • Portable Toilets • Shower Areas • Sinks | • Sports Equipment • Tables • Telephones • Toilet Seats | Soft surfaces | • Bedding • Couches • Curtains • Mattresses |
|----------------------------------|--|--|---|--|----------------------|--|

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

To Disinfect: Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes then allow to air dry. **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry. **Fabric Sanitizer:** To Sanitize Soft Surfaces: Spray until fabric is wet. DO NOT SATURATE. Fabric must remain wet for 30 seconds. Let air dry. For difficult odors, repeat application.

Rinse toys and food contact surfaces with potable water after use.

To Control and Prevent Mold & Mildew: Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals when mold and mildew growth appears. **To Deodorize:** Spray on surfaces as needed. Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

For Tuberculocidal Activity: This product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.

*****KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1 AIDS VIRUS), HEPATITIS B VIRUS (HBV), AND HEPATITIS C VIRUS (HCV) ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids; and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS), Hepatitis B Virus, and Hepatitis C Virus.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV, AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS: PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1, HBV, and HCV. For other organisms, surfaces must remain wet for 3 minutes. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

CONTACT NUMBER: Questions? Comments or In case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

PHYSICAL HAZARDS: FLAMMABLE: Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed. **DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. This can be made from an average of 25% recycled steel (10% post-consumer).

Questions? Call 1-800-228-4722

For ingredient and other information, visit www.thescience.com

Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in U.S.A. © 2016 RB

EPA Reg. No.: 777-99-675

EPA Est. No.: 777-NU-2 (See Bottom for Lot/Date code)

8232786




PLEASE RECYCLE WHEN EMPTY



FOR COMMERCIAL USE

KILLS COLD & FLU VIRUSES†



KILLS 99.9% OF VIRUSES & BACTERIA**

Disinfectant Spray

Eliminates odors

Fresh scent



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. 9833.249

NET WT.:
19 OZ.
(1 LB. 3 OZ.)
538 g

ACTIVE INGREDIENTS:

Alkyl (50% C14, 40% C12, 10% C16)
dimethyl benzyl ammonium saccharinate.....0.10%
Ethanol.....58.00%
OTHER INGREDIENTS.....41.90%
TOTAL.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS.

- Germicidal • Virucidal* • Fungicidal • Tuberculocidal • Bactericidal • Mildewcidal • Kills antibiotic resistant bacteria*
- Kills the source of mold allergens and prevents it from coming back for up to 7 days††
- Deodorizes by killing odor-causing bacteria at their source. Ordinary, non-germicidal sprays can't do this.
- Use on soft surfaces where bacteria can hide*** Meets OSHA Bloodborne Pathogens Standards§§
- Meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants
- ***Kills 99.9% of Staphylococcus aureus and Klebsiella pneumoniae on soft surfaces in 30 seconds

| | | | |
|-----------------------------|--|--|---|
| Kills in 30 seconds: | • Hepatitis C Virus • Herpes Simplex Virus Type 1 • Herpes Simplex Virus Type 2 • Human Coronavirus • Human Immunodeficiency Virus | • Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus* • Influenza B Virus • Newcastle Disease Virus • Respiratory Syncytial Virus (RSV)** | • Rhinovirus Type 39† • Rotavirus WA** • SARS Coronavirus • Swine Influenza A Virus • Vaccinia Virus |
| Kills in 2 minutes: | • Enterovirus Type 68 • Escherichia coli • Haemophilus influenzae • Klebsiella pneumoniae • Klebsiella pneumoniae NDM-1 positive • Klebsiella pneumoniae Carbapenem resistant • Listeria monocytogenes • Neisseria elongata • Pasteurella multocida • Poliovirus Type 1 | • Proteus mirabilis • Proteus vulgaris • Pseudomonas aeruginosa** • Salmonella enterica** • Serratia marcescens • Shigella dysenteriae • Shigella flexneri • Shigella sonnei • Staphylococcus aureus** • Staphylococcus aureus - Methicillin resistant (HA-MRSA)* • Staphylococcus epidermidis | • Staphylococcus haemolyticus • Staphylococcus saprophyticus • Streptococcus pneumoniae PRSP • Streptococcus pyogenes (Strep) • Streptococcus salivarius (Pink Eye) • Trichophyton mentagrophytes (Athlete's Foot Fungus) (Ringworm) • Vibrio cholerae • Yersinia enterocolitica |
| Kills in 3 minutes: | • Adenovirus 14 | • Aspergillus niger† | • Bordetella avium |
| Kills in 10 minutes: | • Feline Calicivirus (Norovirus) | • Mycobacterium bovis BCG (Quant tuberculosis) | |

Intended for use in:

- Ambulances • Clinics • Dental Offices • Hotel/Motel • Office Buildings • Physicians Offices • Schools
- Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices
- Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

| | | | | | | |
|----------------------------------|--|--|---|--|----------------------|--|
| Hard, non-porous surfaces | • Bathroom Surfaces • Chairs • Counters • Desks | • Door Handles • Examination Tables • Floors • Garbage Cans | • Light Switches • Portable Toilets • Shower Areas • Sinks | • Sports Equipment • Tables • Telephones • Toilet Seats | Soft surfaces | • Bedding • Couches • Curtains • Mattresses |
|----------------------------------|--|--|---|--|----------------------|--|

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

To Disinfect: Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes then allow to air dry. **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry. **Fabric Sanitizer:** To Sanitize Soft Surfaces: Spray until fabric is wet. DO NOT SATURATE. Fabric must remain wet for 30 seconds. Let air dry. For difficult odors, repeat application.

Rinse toys and food contact surfaces with potable water after use.

To Control and Prevent Mold & Mildew: Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears. **To Deodorize:** Spray on surfaces as needed.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

For Tuberculocidal Activity: This product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.

***KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1 AIDS VIRUS), HEPATITIS B VIRUS (HBV), AND HEPATITIS C VIRUS (HCV) ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids; and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS), Hepatitis B Virus, and Hepatitis C Virus.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV, AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS: PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1, HBV, and HCV. For other organisms, surfaces must remain wet for 3 minutes. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

CONTACT NUMBER: Questions? Comments or In case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

PHYSICAL HAZARDS: FLAMMABLE: Contents under pressure. Keep away from heat, sparks and open flame.

Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed. **DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. This can be made from an average of 25% recycled steel (10% post-consumer).

Questions? Call 1-800-228-4722

For ingredient and other information, visit www.rbzinfo.com

Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224

Made in U.S.A. © 2016 RB


EPA Reg. No.: 777-99-6175

EPA Est. No.: 777-AU-2 (See Bottom for Lot/Date code) 8232785



FOR COMMERCIAL USE

KILLS COLD & FLU VIRUSES†



KILLS 99.9% OF VIRUSES & BACTERIA**

Disinfectant Spray

Eliminates odors

Original scent



STATE OF HAWAII
Department of Agriculture

ACCEPTED
LICENSE NO. 9833.249

NET WT.:
19 OZ.
(1 LB. 3 OZ.)
538 g

ACTIVE INGREDIENTS:
Alkyl (50% C14, 40% C12, 10% C16) dimethyl benzyl ammonium saccharinate...0.10%
Ethanol...58.00%
OTHER INGREDIENTS:...41.90%
TOTAL:...100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

- Germicidal • Virucidal¹ • Fungicidal • Tuberculocidal • Bactericidal • Mildewcidal • Kills antibiotic resistant bacteria⁴
- Kills the source of mold allergens and prevents it from coming back for up to 7 days¹¹
- Deodorizes by killing odor-causing bacteria at their source. Ordinary, non-germicidal sprays can't do this.
- Use on soft surfaces where bacteria can hide*** Meets OSHA Bloodborne Pathogens Standards³⁸
- Meets ADAC Germicidal Spray Product Test standards for hospital aerosol disinfectants
- ***Kills 99.9% of Staphylococcus aureus and Klebsiella pneumoniae on soft surfaces in 30 seconds

| | | | |
|----------------------------------|--|--|--|
| Kills in 30 seconds: | • Hepatitis C Virus | Type 1 [HIV-1] [AIDS Virus] | • Rhinovirus Type 39 ¹ |
| • Avian Influenza A Virus | • Herpes Simplex Virus Type 1 | • Influenza A Virus ¹ | • Rotavirus WA ¹¹ |
| • Campylobacter jejuni | • Herpes Simplex Virus Type 2 | • Influenza B Virus | • SARS Coronavirus |
| • Coxsackie Type B3 Virus | • Human Coronavirus | • Newcastle Disease Virus | • Swine Influenza A Virus |
| • Cytomegalovirus | • Human Immunodeficiency Virus | • Respiratory Syncytial Virus (RSV) ¹¹ | • Vaccinia Virus |
| • Hepatitis B Virus | | | |
| Kills in 2 minutes: | • Enterovirus Type 68 | • Proteus mirabilis | • Staphylococcus haemolyticus |
| • Acinetobacter baumannii MDR | • Escherichia coli | • Proteus vulgaris | • Staphylococcus saprophyticus |
| • Bordetella pertussis | • Haemophilus influenzae | • Pseudomonas aeruginosa ¹¹ | • Streptococcus pneumoniae PRSP |
| • Burkholderia cepacia | • Klebsiella pneumoniae | • Salmonella enterica ¹¹ | • Streptococcus pyogenes (Strep) |
| • Candida albicans | • Klebsiella pneumoniae NDM-1 positive | • Serratia marcescens | • Streptococcus salivarius (Pink Eye) |
| • Candida parapsilosis | • Klebsiella pneumoniae Carbapenem resistant | • Shigella dysenteriae | • Trichophyton mentagrophytes (Athlete's Foot Fungus) (Ringworm) |
| • Citrobacter freundii | • Klebsiella pneumoniae Listeria monocytogenes | • Shigella flexneri | • Yersinia enterocolitica |
| • Corynebacterium diphtheriae | • Neisseria elongata | • Shigella sonnei | |
| • Cronobacter sakazakii | • Pasteurella multocida | • Staphylococcus aureus ¹¹ | |
| • Enterobacter aerogenes MDR | • Poliovirus Type 1 | • Staphylococcus aureus - Methicillin resistant (HA-MRSA) ¹ | |
| • Enterococcus faecalis | | • Staphylococcus epidermidis | |
| • Enterococcus faecium VRE + MDR | | | |
| Kills in 3 minutes: | • Adenovirus 14 | • Aspergillus niger ¹¹ | • Bordetella avium |
| Kills in 10 minutes: | • Feline Calicivirus (Norovirus) | • Mycobacterium bovis BCG (Quant tuberculosis) | |

Intended for use in:

- Ambulances • Clinics • Dental Offices • Hotel/Motel • Office Buildings • Physicians Offices • Schools
- Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices
- Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

| | | | | | | |
|----------------------------------|----------------------|--------------------|------------------|--------------------|----------------------|------------|
| Hard, non-porous surfaces | • Bathroom Surfaces | • Door Handles | • Light Switches | • Sports Equipment | Soft surfaces | • Bedding |
| • Chairs | • Examination Tables | • Portable Toilets | • Shower Areas | • Tables | • Couches | • Curtains |
| • Counters | • Floors | • Telephone | • Toilet Seats | • Toilets | • Mattresses | |
| • Desks | • Garbage Cans | • Sinks | | | | |

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

To Disinfect: Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes then allow to air dry. **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry. **Fabric Sanitizer:** To Sanitize Soft Surfaces: Spray until fabric is wet. DO NOT SATURATE. Fabric must remain wet for 30 seconds. Let air dry. For difficult odors, repeat application.

Rinse toys and food contact surfaces with potable water after use.

To Control and Prevent Mold & Mildew: Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears. **To Deodorize:** Spray on surfaces as needed. Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

For Tuberculocidal Activity: This product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.

KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1 AIDS VIRUS), HEPATITIS B VIRUS (HBV), AND HEPATITIS C VIRUS (HCV) ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids; and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS), Hepatitis B Virus, and Hepatitis C Virus.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV, AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS: PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1, HBV, and HCV. For other organisms, surfaces must remain wet for 3 minutes. For Norovirus and Mycobacterium bovis BCG surfaces must remain wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

CONTACT NUMBER: Questions? Comments or in case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

PHYSICAL HAZARDS: FLAMMABLE: Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed. **DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. This can be made from an average of 25% recycled steel (10% post-consumer).

Questions? Call 1-800-228-4722
For ingredient and other information, visit www.rbx2info.com


Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in U.S.A. © 2016 RB
EPA Reg. No.: 777-99-675
EPA Est. No.: 777-1U-2 (See Bottom for Lot/Date code) 8232789



PLEASE RECYCLE WHEN EMPTY



FOR COMMERCIAL USE KILLS COLD & FLU VIRUSES[†]



KILLS 99.9% OF VIRUSES & BACTERIA¹¹

Disinfectant Spray

Eliminates odors

Spring Waterfall[®] scent



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. 9833.249

NET WT.:
19 OZ.
(1 LB. 3 OZ.)
538 g

ACTIVE INGREDIENTS:

Alkyl (50% C14, 40% C12, 10% C16)
dimethyl benzyl ammonium saccharinate... 0.10%
Ethanol... 58.00%
OTHER INGREDIENTS... 41.90%
TOTAL... 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS.

This product meets ADAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.

GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL

Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

****Disinfects:** Kills the following household germs.

Kills in 30 seconds:

• Influenza A Virus (New Caledonia/20/99)† • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hepatitis B Virus (HBV) • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) [AIDS Virus] • Influenza A Virus (H1N1)† • Influenza B Virus (Strain B/Hong Kong/572) • Respiratory Syncytial Virus (RSV) • Rhinovirus Type 39† • Rotavirus WA • Vaccinia Virus

Kills in 3 minutes:

• Acinetobacter calcoaceticus • Aspergillus niger (mold & mildew)†† • Burkholderia cepacia • Candida albicans • Corynebacterium diphtheriae • Echovirus Type 12 • Enterobacter aerogenes MDR • Enterococcus faecalis • Enterococcus faecalis - Vancomycin Resistant (VRE) • Escherichia coli (O157:H7) • Klebsiella pneumoniae • Klebsiella pneumoniae NDM-1 positive • Listeria monocytogenes • Neisseria elongata • Penicillium chrysogenum†† • Poliovirus Type 1 • Proteus mirabilis • Proteus vulgaris • Pseudomonas aeruginosa • Pseudomonas putida • Salmonella enterica • Salmonella enterica serovar enteritidis • Salmonella enterica serovar typhi • Serratia marcescens • Shigella dysenteriae • Staphylococcus aureus • Staphylococcus aureus - Methicillin Resistant (HA-MRSA) • Staphylococcus epidermidis • Streptococcus pyogenes (Strep) • Streptococcus salivarius (Pink Eye) • Trichophyton mentagrophytes (Athlete's Foot Fungus) • Klebsiella pneumoniae Carbapenem Resistant

Kills in 10 minutes:

BACTERIA:

• Mycobacterium bovis BCG (Quant tuberculosis)

VIRUSES:

• Feline calicivirus (Norovirus)

Intended for use only in:

• Ambulances • Clinics • Dental Offices • Hotel/Motel • Nurseries • Physicians Offices • Schools
• Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices
• Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces • Door handles • Floors • Patient Chairs • Shower Areas • Telephones
• Bed Frames • Dressing carts • Faucets • Plastic Toys† • Sinks • Window Sills
• Counters • Drinking Fountains • Glazed Porcelain • Portable Toilets • Sports Equipment • Wheelchairs
• Changing Tables • Examination tables • Linen carts • Salad Bar Sneeze Guards • Stainless Steel* • Whirlpool Interiors

† Follow directions For Toys under Directions for Use - rinsing application required. * Non-Medical or Fixture

This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.

KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) [AIDS VIRUS] AND Hepatitis B Virus (HBV) ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and HBV. For other organisms, surfaces must remain wet for 3 minutes. For Norovirus and Mycobacterium bovis BCG (Quant tuberculosis) surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

- **To Deodorize:** Spray on surfaces as needed.
- **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry.
- **To Disinfect:** Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and Mycobacterium bovis BCG (Quant tuberculosis) surfaces must remain wet for 10 minutes then allow to air dry.
- **Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
• Call a Poison Control Center or doctor for treatment advice.

CONTACT NUMBER

Questions? Comments or In case of an emergency, call toll free 1-800-228-4722.
Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

PHYSICAL HAZARDS: FLAMMABLE. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

DISPOSAL: Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

Questions? Call 1-800-228-4722

For ingredient and other information, www.rbnainfo.com

Distributed by: Reckitt Benckiser
Parsippany, NJ 07054-0224
Made in U.S.A. © 2016 RB

EPA Reg. No.: 777-99-675

EPA Est. No.: 777-NJ-2

(See Bottom for Lot/Date code)

8223292



HEALTH-HYGIENE-HOME




PLEASE RECYCLE
WHEN EMPTY



LISTED (139865)

KILLS COLD & FLU VIRUSES†



Disinfectant Spray

Eliminates odors

Crisp Linen®
scent



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. 9833.249

NET WT.:
19 OZ.
(1 LB. 3 OZ.)
538 g

ACTIVE INGREDIENTS:

Alkyl (50% C14, 40% C12, 10% C16)
dimethyl benzyl ammonium saccharinate...0.10%
Ethanol.....58.00%

OTHER INGREDIENTS:.....41.90%

TOTAL:.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

This product meets ADAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.
GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL*
Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

****Disinfects:** Kills the following household germs.

Kills in 30 seconds:

• Influenza A Virus (New Caledonia/20/99)† • *Campylobacter jejuni* • *Coxsackie* Type B3 Virus • *Orytomegalovirus* • *Hepatitis B Virus* (HBV) • *Herpes Simplex Virus* Type 1 & 2 • *Human Immunodeficiency Virus* Type 1 (HIV-1) (AIDS Virus) • *Influenza A Virus* (H1N1)† • *Influenza B Virus* (Strain B/Hong Kong/572) • *Respiratory Syncytial Virus* (RSV) • *Rhinovirus* Type 39† • *Rotavirus* WA • *Vaccinia Virus*

Kills in 3 minutes:

• *Aspergillus niger* (mold & mildew)†† • *Burkholderia cepacia* • *Candida albicans* • *Corynebacterium diphtheriae* • *Echovirus* Type 12
• *Enterobacter aerogenes* MD8 • *Enterococcus faecalis* • *Enterococcus faecalis* - Vancomycin Resistant (VRE) • *Escherichia coli* (O157:H7) • *Klebsiella pneumoniae*
• *Klebsiella pneumoniae* ND8M-1 positive • *Listeria monocytogenes* • *Neisseria elongata* • *Penicillium chrysogenum*†† • *Poliovirus* Type 1 • *Proteus mirabilis* • *Proteus vulgaris* • *Pseudomonas aeruginosa* • *Pseudomonas putida* • *Salmonella enterica* • *Salmonella enterica* serovar enteritidis • *Salmonella enterica* serovar typhi • *Serratia marcescens* • *Shigella dysenteriae* • *Staphylococcus aureus* • *Staphylococcus aureus* - Methicillin Resistant (HA-MRSA) • *Staphylococcus epidermidis* • *Streptococcus pyogenes* (Strep) • *Streptococcus salivarius* (Pink Eye) • *Trichophyton mentagrophytes* (Athlete's Foot Fungus) • *Klebsiella pneumoniae* Carbapenem Resistant

Kills in 10 minutes:

BACTERIA:

• *Mycobacterium bovis* BCG (Quant tuberculosis)

VIRUSES:

• *Feline calicivirus* (Norovirus)

Intended for use only in:

• Ambulances • Clinics • Dental Offices • Hotel/Motel • Nurseries • Physicians Offices • Schools
• Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices
• Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces • Door handles • Floors • Patient Chairs • Shower Areas • Telephones
• Bed Frames • Dressing carts • Faucets • Plastic Toys† • Sinks • Windowsills
• Counters • Drinking Fountains • Glazed Porcelain • Portable Toilets • Sports Equipment • Wheelchairs
• Changing Tables • Examination tables • Linen carts • Salad Bar Sneeze Guards • Stainless Steel† • Whirlpool Interiors

† Follow directions For Toys under Directions for Use - rinsing application required. †† Non-Medical or Fixture

This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.

KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) AND Hepatitis B Virus (HBV) ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids; and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

• **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.

• **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

• **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and HBV. For other organisms, surfaces must remain wet for 3 minutes.

For Norovirus and *Mycobacterium bovis* BCG (Quant tuberculosis) surfaces must remain wet for 10 minutes.

• **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

- **To Deodorize:** Spray on surfaces as needed.
- **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry.
- **To Disinfect:** Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and *Mycobacterium bovis* BCG (Quant tuberculosis) surfaces must remain wet for 10 minutes then allow to air dry.
- **Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
• Call a Poison Control Center or doctor for treatment advice.

CONTACT NUMBER

Questions? Comments or In case of an emergency, call toll free 1-800-228-4722.
Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

PHYSICAL HAZARDS: FLAMMABLE. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

DISPOSAL: Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

Questions? (1-800-228-4722

For ingredient and other information, www.rbnainfo.com


Distributed by: Reckitt Benckiser
Parsippany, NJ 07054-0224
Made in U.S.A. © 2016 RB

EPA Reg. No.: 777-99-675
EPA Est. No.: 777-NJ-2
(See Bottom for Lot/Date code)

8223288



KILLS COLD & FLU VIRUSES*



Disinfectant Spray

Eliminates odors

Early Morning Breeze scent



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. 9833.249

NET WT.:
19 OZ.
(1 LB. 3 OZ.)
538 g

ACTIVE INGREDIENTS:

Alkyl (50% C14, 40% C12, 10% C16)
dimethyl benzyl ammonium saccharinate...0.10%
Ethanol.....58.00%
OTHER INGREDIENTS:.....41.90%
TOTAL:.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

8 BOX TOPS



STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO. 9833.249



KILLS COLD & FLU VIRUSES†

4 PACK

**Disinfectant
Spray**
Eliminates odors

Crisp Linen® scent

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

NET WT. 4 x 19 OZ. (1 LB. 3 OZ.) 538 g
TOTAL WT. 76 OZ. (4 LBS. 12 OZ.) 2.1 kg

ACTIVE INGREDIENTS:
Alkyl (50% C14, 40% C12, 10% C16).....0.10%
dimethyl benzyl ammonium saccharinate...0.10%
Ethanol.....58.00%
OTHER INGREDIENTS.....41.90%
TOTAL.....100.00%



This product meets ADAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.
GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL†

Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

****Disinfects:** Kills the following household germs.

Kills in 30 seconds:

- Influenza A Virus (New Caledonia/20/99)† • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hepatitis B Virus (HBV) • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus (H1N1)† • Influenza B Virus (Strain B/Hong Kong/5/72) • Respiratory Syncytial Virus (RSV) • Rhinovirus Type 391 • Rotavirus WA • Vaccinia Virus

Kills in 3 minutes:

- Acinetobacter calcoaceticus • Aspergillus niger (mold & mildew)†† • Burkholderia cepacia • Candida albicans • Corynebacterium diphtheriae • Echovirus Type 12 • Enterobacter aerogenes MDR • Enterococcus faecalis • Enterococcus faecalis - Vancomycin Resistant (VRE) • Escherichia coli (O157:H7) • Klebsiella pneumoniae • Klebsiella pneumoniae - NDM-1 positive • Listeria monocytogenes • Neisseria elongata • Penicillium chrysogenum†† • Poliovirus Type 1 • Proteus mirabilis • Proteus vulgaris • Pseudomonas aeruginosa • Pseudomonas putida • Salmonella enterica • Salmonella enterica serovar enteritidis • Salmonella enterica serovar typhi • Serratia marcescens • Shigella dysenteriae • Staphylococcus aureus • Staphylococcus aureus - Methicillin Resistant (HA-MRSA) • Staphylococcus epidermidis • Streptococcus pyogenes (Strep) • Streptococcus salivarius (Pink Eye) • Trichophyton mentagrophytes (Athlete's Foot Fungus) • Klebsiella pneumoniae Carbapenem Resistant

Kills in 10 minutes:

BACTERIA:

- Mycobacterium bovis BCG (Quant tuberculosis)

VIRUSES:

- Feline calicivirus (Norovirus)

Intended for use only in:

- Ambulances • Clinics • Dental Offices • Hotel/Motel • Nurseries • Physicians Offices • Schools
- Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Restrooms • Veterinary Offices
- Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

- Bathroom surfaces • Door handles • Floors • Patient Chairs • Shower Areas • Telephones
- Bed Frames • Dressing carts • Faucets • Plastic Toys • Sinks • Windowsills
- Counters • Drinking Fountains • Glazed Porcelain • Portable Toilets • Sports Equipment • Wheelchairs
- Changing Tables • Examination tables • Linen carts • Salad Bar Sneeze Guards • Stainless Steel* • Whirlpool Interiors

† Follow directions For Toys under Directions for Use - rinsing application required. *Non-Medical or Fixture

This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.

KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) AND HEPATITIS B VIRUS (HBV) ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids; and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and HBV. For other organisms, surfaces must remain wet for 3 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Preclean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

- **To Deodorize:** Spray on surfaces as needed.
- **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry.
- **To Disinfect:** Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and Mycobacterium bovis BCG (Quant tuberculosis) surfaces must remain wet for 10 minutes then allow to air dry.
- **Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. **Do not** spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

| FIRST AID | |
|---|--|
| If in eyes | • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice. |
| CONTACT NUMBER | |
| Questions? Comments or in case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment. | |

PHYSICAL HAZARDS: FLAMMABLE. Contents under pressure. Keep away from heat, sparks and open flame.

Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal.

STORAGE: Store in original container in areas inaccessible to children. Keep securely closed.

DISPOSAL: Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

Questions? Call 1-800-228-4722

For ingredient and other information, www.tbainfo.com

Distributed by: Reckitt Benckiser

Parsippany, NJ 07054-1224

Made in U.S.A. © 2016 RB

EPA Reg. No.: 777-99-675

EPA Est. No.: 777-NJ-2

(See Bottom for Lot/Date code)

8254838



Do Not Remove label/cut out Box Tops for Education proof of purchase until product has been completely used. Box Tops for Education and associated words and designs are trademarks of General Mills, used under license. © General Mills

Go to boxtops4education.com for details



This product meets ADAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.

GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL†

Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

****Disinfects:** Kills the following household germs.

Kills in 30 seconds:

• Influenza A Virus (New Caledonia/20/99)† • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hepatitis B Virus (HBV) • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) [AIDS Virus] • Influenza A Virus (H1N1)† • Influenza B Virus (Strain B/Hong Kong/572) • Respiratory Syncytial Virus (RSV) • Rhinovirus Type 39† • Rotavirus WA • Vaccinia Virus

Kills in 3 minutes:

• Acinetobacter calcoaceticus • Aspergillus niger (mold & mildew)†† • Burkholderia cepacia • Candida albicans • Corynebacterium diphtheriae • Echovirus Type 12 • Enterobacter aerogenes MDR • Enterococcus faecalis • Enterococcus faecalis - Vancomycin Resistant (VRE) • Escherichia coli (O157:H7) • Klebsiella pneumoniae • Klebsiella pneumoniae NDM-1 positive • Listeria monocytogenes • Neisseria elongata • Penicillium chrysogenum†† • Poliovirus Type 1 • Proteus mirabilis • Proteus vulgaris • Pseudomonas aeruginosa • Pseudomonas putida • Salmonella enterica • Salmonella enterica serovar enteritidis • Salmonella enterica serovar typhi • Serratia marcescens • Shigella dysenteriae • Staphylococcus aureus • Staphylococcus aureus - Methicillin Resistant (HA-MRSA) • Staphylococcus epidermidis • Streptococcus pyogenes (Strep) • Streptococcus salivarius (Pink Eye) • Trichophyton mentagrophytes (Athlete's Foot Fungus) • Klebsiella pneumoniae Carbapenem Resistant

Kills in 10 minutes:

BACTERIA:

• Mycobacterium bovis BCG (Quant tuberculosis)

VIRUSES:

• Feline calicivirus (Norovirus)

Intended for use only in:

• Ambulances • Clinics • Dental Offices • Hotel/Motel • Nurseries • Physicians Offices • Schools
• Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices
• Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces • Door handles • Floors • Patient Chairs • Shower Areas • Telephones
• Bed Frames • Dressing carts • Faucets • Plastic Toys† • Sinks • Window Sills
• Counters • Drinking Fountains • Glazed Porcelain • Portable Toilets • Sports Equipment • Wheelchairs
• Changing Tables • Examination tables • Linen carts • Salad Bar Sneeze Guards • Stainless Steel* • Whirlpool Interiors

† Follow directions For Toys under Directions for Use - rinsing application required. * Non-Medical or Fixture

This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.

KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) [AIDS VIRUS] AND Hepatitis B Virus (HBV) ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids; and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and HBV. For other organisms, surfaces must remain wet for 3 minutes. For Norovirus and Mycobacterium bovis BCG (Quant tuberculosis) surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 3 to 4 seconds until covered with mist.

- **To Deodorize:** Spray on surfaces as needed.
- **To Sanitize:** Surfaces must remain wet for 10 seconds then allow to air dry.
- **To Disinfect:** Surfaces must remain wet for 3 minutes then allow to air dry. For Norovirus and Mycobacterium bovis BCG (Quant tuberculosis) surfaces must remain wet for 10 minutes then allow to air dry.
- **Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
• Call a Poison Control Center or doctor for treatment advice.

CONTACT NUMBER

Questions? Comments or In case of an emergency, call toll free 1-800-228-4722.
Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

PHYSICAL HAZARDS: FLAMMABLE. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

DISPOSAL: Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

Questions? Call 1-800-228-4722

For ingredient and other information, www.rbnainfo.com

Distributed by: Reckitt Benckiser
Parsippany, NJ 07054-0224
Made in U.S.A. © 2016 RB

EPA Reg. No.: 777-99-675

EPA Est. No.: 777-NJ-2

(See Bottom for Lot/Date code)

8223292



HEALTH & HYGIENE - HOME



PLEASE RECYCLE
WHEN EMPTY



KILLS COLD & FLU VIRUSES†

Professional Lysol®

BRAND III

KILLS 99.9% OF VIRUSES & BACTERIA**

Disinfectant Spray

Eliminates odors

Crisp Linen®
scent

NET WT.:
19 OZ.
(1 LB. 3 OZ.)
538 g

ACTIVE INGREDIENTS:

Alkyl (50% C14, 40% C12, 10% C16)
dimethyl benzyl ammonium saccharinate...0.10%
Ethanol.....58.00%

OTHER INGREDIENTS:.....41.90%

TOTAL:.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS