

This product meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.

GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL†

Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

**Kills in 30 seconds:**

• Avian Influenza A Virus • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hantavirus • Hepatitis B Virus • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus† • Influenza B Virus • Respiratory Syncytial Virus • Rhinovirus Type 39† • Rotavirus WA • Vaccinia Virus

**Kills in 5 minutes:**

• Enterobacter aerogenes • Vancomycin Resistant Enterococcus faecalis (VRE) • Escherichia coli (O157:H7) • Listeria monocytogenes • Pseudomonas aeruginosa • Salmonella enterica • Staphylococcus aureus • Trichophyton mentagrophytes (Athlete's Foot Fungus)

**\*\*Kills in 10 minutes:**

**BACTERIA:**

• Acinetobacter calcoaceticus	• Klebsiella pneumoniae	• Proteus mirabilis	• Shigella dysenteriae
• Burkholderia cepacia	• Carbapenem Resistant	• Proteus vulgaris	• Staphylococcus epidermidis
• Corynebacterium diphtheriae	• Methicillin Resistant	• Pseudomonas putida	• Streptococcus pyogenes
• Enterococcus faecalis	• Staphylococcus aureus (MRSA)	• Salmonella enterica serovar enteritidis	• Streptococcus salivarius
• Klebsiella pneumoniae	• Mycobacterium bovis BCG (Quant tuberculosis)	• Salmonella enterica serovar Paratyphi B	
• Klebsiella pneumoniae - NDM-1 positive	• Neisseria elongata	• Salmonella enterica serovar typhi	
		• Serratia marcescens	

**FUNGI:**

• Aspergillus niger (mold & mildew)†† • Candida albicans • Fusarium solani • Penicillium chrysogenum††

**VIRUSES:**

• Adenovirus Type 2 • Echovirus Type 12 • Hepatitis A Virus • Poliovirus Type 1 • Feline calicivirus (Norovirus)

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces	• Door handles	• Floors	• Patient Chairs	• Shower Areas	• Telephones
• Bed Frames	• Dressing carts	• Faucets	• Plastic Toys†	• Sinks	• Windowills
• Counters	• Drinking Fountains	• Glazed Porcelain	• Portable Toilets	• Sports Equipment	• Wheelchairs
• Changing Tables	• Examination tables	• Linen carts	• Salad Bar Sneeze Guards	• Stainless Steel†	• Whirlpool Interiors

† Follow directions For Toys under Directions for Use - rinsing application required. †† Non-Medical or Fixture

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) AND HEPATITIS B VIRUS ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HEPATITIS B VIRUS ON SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS:**

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and Hepatitis B Virus. For all other organisms, surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

Intended for use only in:

• Ambulances	• Clinics	• Dental Offices	• Hotel/Motel	• Nurseries	• Physicians Offices	• Schools
• Blood Banks	• Colleges	• Health Clubs	• Laboratories	• Nursing Homes	• Restaurants	• Waiting Offices
• Cafeterias	• Day Care Centers	• Hospitals	• Military Installations	• Pharmacies	• Restrooms	• Veterinary Rooms

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. This product is not for use on critical & semi-critical medical devices or medical equipment surfaces. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 2 to 3 seconds until covered with mist.

- **To Disinfect:** Let stand for 10 minutes then allow to air dry.
- **Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Sanitize:** Let stand for 30 seconds then allow to air dry.
- **To Deodorize:** Spray on surfaces as needed.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

**PRECAUTIONARY STATEMENTS:** Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
<b>If in eyes</b>	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
CONTACT NUMBER	
Questions? Comments or in case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.	

**PHYSICAL HAZARDS: FLAMMABLE.** Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

**DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

**Questions? (1-800-228-4722)**

For ingredient and other information, [www.rbninfo.com](http://www.rbninfo.com)

Distributed by: Reckitt Benckiser  
Parsippany, NJ 07054-0224  
Made in U.S.A. © 2012 RB

EPA Reg. No.: 777-99-675  
EPA Est. No.: 777-NJ-2  
(See Bottom for Lot/Date code)

0396306



Federal regulations prohibit CFC propellants in aerosols.



PLEASE RECYCLE WHEN EMPTY



0 36241 76075 5

**KILLS COLD & FLU VIRUSES\***



**KILLS 99.9% OF VIRUSES & BACTERIA\*\***

**Disinfectant Spray**

**Spring Waterfall® scent**



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. **9833.249**

NET WT.:  
19 OZ.  
(1 LB. 3 OZ.)  
538 g

**ACTIVE INGREDIENTS:**  
Alkyl (50% C14, 40% C12, 10% C16) dimethyl benzyl ammonium saccharinate...0.10%  
Ethanol...58.00%  
**OTHER INGREDIENTS:**...41.90%  
**TOTAL:**...100.00%

KEEP OUT OF REACH OF CHILDREN  
**CAUTION:** SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

This product meets AOC Germicidal Spray Product Test standards for hospital aerosol disinfectants.

GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL†

Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

**Kills in 30 seconds:**

• Avian Influenza A Virus • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hantavirus • Hepatitis B Virus • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus† • Influenza B Virus • Respiratory Syncytial Virus • Rhinovirus Type 39F • Rotavirus WA • Vaccinia Virus

**Kills in 5 minutes:**

• Enterobacter aerogenes • Vancomycin Resistant Enterococcus faecalis (VRE) • Escherichia coli (O157:H7) • Listeria monocytogenes • Pseudomonas aeruginosa • Salmonella enterica • Staphylococcus aureus • Trichophyton mentagrophytes (Athlete's Foot Fungus)

**\*\*Kills in 10 minutes:**

**BACTERIA:**

• Acinetobacter calcoaceticus	• Klebsiella pneumoniae	• Proteus mirabilis	• Shigella dysenteriae
• Burkholderia cepacia	• Carbapenem Resistant	• Proteus vulgaris	• Staphylococcus epidermidis
• Corynebacterium diphtheriae	• Methicillin Resistant	• Pseudomonas putida	• Streptococcus pyogenes
• Enterococcus faecalis	• Staphylococcus aureus (MRSA)	• Salmonella enterica serovar enteritidis	• Streptococcus salivarius
• Klebsiella pneumoniae	• Mycobacterium bovis BCG	• Salmonella enterica serovar Paratyphi B	
• Klebsiella pneumoniae	• (Quant tuberculosis)	• Salmonella enterica serovar typhi	
• NDM-1 positive	• Neisseria elongata	• Serratia marcescens	

**FUNGI:**

• Aspergillus niger (mold & mildew)†† • Candida albicans • Fusarium solani • Penicillium chrysogenum††

**VIRUSES:**

• Adenovirus Type 2 • Echovirus Type 12 • Hepatitis A Virus • Poliovirus Type 1 • Feline calicivirus (Norovirus)

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces	• Door handles	• Floors	• Patient Chairs	• Shower Areas	• Telephones
• Bed Frames	• Dressing carts	• Faucets	• Plastic Toys	• Sinks	• Windowills
• Counters	• Drinking Fountains	• Glazed Porcelain	• Portable Toilets	• Sports Equipment	• Wheelchairs
• Changing Tables	• Examination tables	• Linen carts	• Salad Bar Sneeze Guards	• Stainless Steel†	• Whirlpool Interiors

† Follow directions for Toys under Directions for Use - rinsing application required. † Non-Medical or Fixture.

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) AND HEPATITIS B VIRUS ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HEPATITIS B VIRUS ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:**

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and Hepatitis B Virus. For all other organisms, surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

**Intended for use only in:**

• Ambulances	• Clinics	• Hotel/Motel	• Nurseries	• Physicians Offices	• Schools
• Blood Banks	• Colleges	• Laboratories	• Nursing Homes	• Restaurants	• Veterinary Offices
• Cafeterias	• Day Care Centers	• Hospitals	• Military Installations	• Restrooms	• Waiting Rooms

**DIRECTIONS FOR USE:** Is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. **This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.** Preclean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 2 to 3 seconds until covered with mist.

- **To Disinfect:** Let stand for 10 minutes then allow to air dry.
- **Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Sanitize:** Let stand for 30 seconds then allow to air dry.
- **To Deodorize:** Spray on surfaces as needed.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

**PRECAUTIONARY STATEMENTS:** Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
<b>If in eyes</b>	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
CONTACT NUMBER	
Questions? Comments or in case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.	

**PHYSICAL HAZARDS:** FLAMMABLE. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed. **DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer). **Questions? (1-800-228-4722)** For ingredient and other information, [www.rbnainfo.com](http://www.rbnainfo.com)

Distributed by: Reckitt Benckiser  
Parsippany, NJ 07054-0224  
Made in U.S.A. © 2012 RB

EPA Reg. No.: 777-99-675  
EPA Est. No.: 777-NJ-2  
(See Bottom for Lot/Date code)  
0396297



Federal regulations prohibit CFC propellants in aerosols.




PLEASE RECYCLE WHEN EMPTY



0 36241 04650 7



**KILLS COLD & FLU VIRUSES†**



**KILLS 99.9% OF VIRUSES & BACTERIA\*\***

**Disinfectant Spray**

**Eliminates odors**

**Original scent**

STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. **9833.249**

NET WT.:  
19 OZ.  
(1 LB. 3 OZ.)  
538 g

**ACTIVE INGREDIENTS:**  
Allyl (50% C14, 40% C12, 10% C16)  
dimethyl benzyl ammonium saccharinate... 0.10%  
Ethanol... 58.00%  
**OTHER INGREDIENTS:**... 41.90%  
**TOTAL:**... 100.00%

KEEP OUT OF REACH OF CHILDREN  
**CAUTION:** SEE BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS

This product meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.  
GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL  
Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

**Kills in 30 seconds:**

• Avian Influenza A Virus • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hantavirus • Hepatitis B Virus • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 [HIV-1] [AIDS Virus] • Influenza A Virus† • Influenza B Virus • Respiratory Syncytial Virus • Rhinovirus Type 39† • Rotavirus WA • Vaccinia Virus

**Kills in 5 minutes:**

• Enterobacter aerogenes • Vancomycin Resistant Enterococcus faecalis (VRE) • Escherichia coli (O157:H7) • Listeria monocytogenes • Pseudomonas aeruginosa • Salmonella enterica • Staphylococcus aureus • Trichophyton mentagrophytes (Athlete's Foot Fungus)

**\*\*Kills in 10 minutes:**

**BACTERIA:**

• Acinetobacter calcoaceticus	• Klebsiella pneumoniae	• Proteus mirabilis	• Shigella dysenteriae
• Burkholderia cepacia	• Carbapenem Resistant	• Proteus vulgaris	• Staphylococcus epidermidis
• Corynebacterium diphtheriae	• Methicillin Resistant	• Pseudomonas putida	• Streptococcus pyogenes
• Enterococcus faecalis	• Staphylococcus aureus (MRSA)	• Salmonella enterica serovar enteritidis	• Streptococcus salivarius
• Klebsiella pneumoniae	• Mycobacterium bovis BCG	• Salmonella enterica serovar Paratyphi B	
• Klebsiella pneumoniae	(Quant tuberculosis)	• Salmonella enterica serovar typhi	
• NDM-1 positive	• Neisseria elongata	• Serratia marcescens	

**FUNGI:**

• Aspergillus niger (mold & mildew)†† • Candida albicans • Fusarium solani • Penicillium chrysogenum††

**VIRUSES:**

• Adenovirus Type 2 • Echovirus Type 12 • Hepatitis A Virus • Poliovirus Type 1 • Feline calicivirus (Norovirus)

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces	• Door handles	• Floors	• Patient Chairs	• Shower Areas	• Telephones
• Bed Frames	• Dressing carts	• Faucets	• Plastic Toys <sup>1</sup>	• Sinks	• Windowills
• Counters	• Drinking Fountains	• Glazed Porcelain	• Portable Toilets	• Sports Equipment	• Wheelchairs
• Changing Tables	• Examination tables	• Linen carts	• Salad Bar Sneeze Guards	• Stainless Steel <sup>1A</sup>	• Whirlpool Interiors

<sup>1</sup> Follow directions for Toys under Directions for Use - rinsing application required. <sup>1A</sup> Non-Medical or Fixture

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) [AIDS VIRUS] AND HEPATITIS B VIRUS ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HEPATITIS B VIRUS ON SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS:**

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and Hepatitis B Virus. For all other organisms, surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

**Intended for use only in:**

• Ambulances	• Clinics	• Dental Offices	• Hotel/Motel	• Nurseries	• Physicians Offices	• Schools
• Blood Banks	• Colleges	• Health Clubs	• Laboratories	• Nursing Homes	• Restaurants	• Veterinary Offices
• Cafeterias	• Day Care Centers	• Hospitals	• Military Installations	• Pharmacies	• Restrooms	• Waiting Rooms

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. **This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.** Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 2 to 3 seconds until covered with mist.

- **To Disinfect:** Let stand for 10 minutes then allow to air dry.
- **Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Sanitize:** Let stand for 30 seconds then allow to air dry.
- **To Deodorize:** Spray on surfaces as needed.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

**PRECAUTIONARY STATEMENTS:** Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. **Do not** spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**FIRST AID**

**If in eyes** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.  
• Call a Poison Control Center or doctor for treatment advice.

**CONTACT NUMBER**

Questions? Comments or in case of an emergency, call toll free 1-800-228-4722.  
Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

**PHYSICAL HAZARDS:** FLAMMABLE. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

**DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

**Questions? (1-800-228-4722)**

For ingredient and other information, [www.rbnainfo.com](http://www.rbnainfo.com)

Distributed by: Reckitt Benckiser  
Parsippany, NJ 07054-0224  
Made in U.S.A. © 2012 RB

EPA Reg. No.: 777-99-675

EPA Est. No.: 777-AU-2

(See Bottom for Lot/Date code)  
0396300




Federal regulations prohibit CFC propellants in aerosols.



PLEASE RECYCLE WHEN EMPTY



**KILLS COLD & FLU VIRUSES\***



**KILLS 99.9% OF VIRUSES & BACTERIA\*\***

**Disinfectant Spray**

**Eliminates odors**

**Fresh scent**



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. **9833.249**

NET WT.:  
19 OZ.  
(1 LB. 3 OZ.)  
538 g

**ACTIVE INGREDIENTS:**

Alkyl (50% C14, 40% C12, 10% C16)	
dimethyl benzyl ammonium saccharinate...	10%
Ethanol.....	58.00%
<b>OTHER INGREDIENTS:</b>	41.90%
<b>TOTAL:</b>	100.00%

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

This product meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.  
GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL\*

Kills the source of mold allergens.†

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

**Kills in 30 seconds:**

• Avian Influenza A Virus • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hantavirus • Hepatitis B Virus • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus • Influenza B Virus • Respiratory Syncytial Virus • Rhinovirus Type 39† • Rotavirus WA • Vaccinia Virus

**Kills in 5 minutes:**

• Enterobacter aerogenes • Vancomycin Resistant Enterococcus faecalis (VRE) • Escherichia coli (O157:H7) • Listeria monocytogenes • Pseudomonas aeruginosa • Salmonella enterica • Staphylococcus aureus • Trichophyton mentagrophytes (Athlete's Foot Fungus)

**\*\*Kills in 10 minutes:**

**BACTERIA:**

• Acinetobacter calcoaceticus	• Klebsiella pneumoniae	• Proteus mirabilis	• Shigella dysenteriae
• Burkholderia cepacia	• Carbapenem Resistant	• Proteus vulgaris	• Staphylococcus epidermidis
• Corynebacterium diphtheriae	• Methicillin Resistant	• Pseudomonas putida	• Streptococcus pyogenes
• Enterococcus faecalis	• Staphylococcus aureus (MRSA)	• Salmonella enterica serovar enteritidis	• Streptococcus salivarius
• Klebsiella pneumoniae	• Mycobacterium bovis BOG (Quart tuberculosis)	• Salmonella enterica serovar Paratyphi B	
• Klebsiella pneumoniae - NDM-1 positive	• Neisseria elongata	• Salmonella enterica serovar typhi	
		• Serratia marcescens	

**FUNGI:**

• Aspergillus niger (mold & mildew)†† • Candida albicans • Fusarium solani • Penicillium chrysogenum††

**VIRUSES:**

• Adenovirus Type 2 • Echovirus Type 12 • Hepatitis A Virus • Poliovirus Type 1 • Feline calicivirus (Norovirus)

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces	• Door handles	• Floors	• Patient Chairs	• Shower Areas	• Telephones
• Bed Frames	• Dressing carts	• Faucets	• Plastic Toys†	• Sinks	• Window Sills
• Counters	• Drinking Fountains	• Glazed Porcelain	• Portable Toilets	• Sports Equipment	• Wheelchairs
• Changing Tables	• Examination tables	• Linen carts	• Salad Bar Sneeze Guards	• Stainless Steel*	• Whirlpool Interiors

† Follow directions For Toys under Directions for Use - rinsing application required. †† Non-Medical or Fixture

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) AND HEPATITIS B VIRUS ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HEPATITIS B VIRUS ON SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS:**

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and Hepatitis B Virus. For all other organisms, surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

Intended for use only in:

• Ambulances	• Clinics	• Dental Offices	• Hotel/Motel	• Nurseries	• Physicians Offices	• Schools
• Blood Banks	• Colleges	• Health Clubs	• Laboratories	• Nursing Homes	• Restaurants	• Veterinary Offices
• Cafeterias	• Day Care Centers	• Hospitals	• Military Installations	• Pharmacies	• Restrooms	• Waiting Rooms

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. **This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.** Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 2 to 3 seconds until covered with mist.

- **To Disinfect:** Let stand for 10 minutes then allow to air dry.
- **To Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Sanitize:** Let stand for 30 seconds then allow to air dry.
- **To Deodorize:** Spray on surfaces as needed.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

**PRECAUTIONARY STATEMENTS:** Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. **Do not** spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**FIRST AID**

<b>If in eyes</b>	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
-------------------	--

**CONTACT NUMBER**

Questions? Comments or in case of an emergency, call toll free 1-800-228-4722.  
Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

**PHYSICAL HAZARDS: FLAMMABLE.** Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

**DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

**Questions? (1-800-228-4722)**

For ingredient and other information, [www.rbnainfo.com](http://www.rbnainfo.com)

Distributed by: Reckitt Benckiser  
Parsippany, NJ 07054-0224  
Made in U.S.A. © 2012 RB

EPA Reg. No.: 777-99-675  
EPA Est. No.: 777-NJ-2  
(See Bottom for Lot/Date code)  
0389720B



Federal regulations prohibit CFC propellants in aerosols.



PLEASE RECYCLE WHEN EMPTY



**KILLS COLD & FLU VIRUSES\***



**KILLS 99.9% OF VIRUSES & BACTERIA\*\***

**Disinfectant Spray**

**Eliminates odors**

**Early Morning Breeze scent**



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. **9833.249**

NET WT.:  
19 OZ.  
(1 LB. 3 OZ.)  
538 g

**ACTIVE INGREDIENTS:**  
Alkyl (50% C14, 40% C12, 10% C16)  
dimethyl benzyl ammonium saccharinate...0.10%  
Ethanol.....58.00%  
**OTHER INGREDIENTS:**.....41.90%  
**TOTAL:**.....100.00%

KEEP OUT OF REACH OF CHILDREN  
**CAUTION:** SEE BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS

This product meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.

GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL†

Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

**Kills in 30 seconds:**

• Avian Influenza A Virus • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hantavirus • Hepatitis B Virus • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus† • Influenza B Virus • Respiratory Syncytial Virus • Rhinovirus Type 39† • Rotavirus WA • Vaccinia Virus

**Kills in 5 minutes:**

• Enterobacter aerogenes • Vancomycin Resistant Enterococcus faecalis (VRE) • Escherichia coli (O157:H7) • Listeria monocytogenes • Pseudomonas aeruginosa • Salmonella enterica • Staphylococcus aureus • Trichophyton mentagrophytes (Athlete's Foot Fungus)

**\*\*Kills in 10 minutes:**

**BACTERIA:**

• Acinetobacter calcoaceticus	• Klebsiella pneumoniae	• Proteus mirabilis	• Shigella dysenteriae
• Burkholderia cepacia	• Carbapenem Resistant	• Proteus vulgaris	• Staphylococcus epidermidis
• Corynebacterium diptheriae	• Methicillin Resistant	• Pseudomonas putida	• Streptococcus pyogenes
• Enterococcus faecalis	• Staphylococcus aureus (MRSA)	• Salmonella enterica serovar enteritidis	• Streptococcus salivarius
• Klebsiella pneumoniae	• Mycobacterium bovis BOG	• Salmonella enterica serovar Paratyphi B	
• Klebsiella pneumoniae - NDM-1 positive	• (Quant tuberculosis)	• Salmonella enterica serovar typhi	
	• Neisseria elongata	• Serratia marcescens	

**FUNGI:**

• Aspergillus niger (mold & mildew)†† • Candida albicans • Fusarium solani • Penicillium chrysogenum††

**VIRUSES:**

• Adenovirus Type 2 • Echovirus Type 12 • Hepatitis A Virus • Poliovirus Type 1 • Feline calicivirus (Norovirus)

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces	• Door handles	• Floors	• Patient Chairs	• Shower Areas	• Telephones
• Bed Frames	• Dressing carts	• Faucets	• Plastic Toys	• Sinks	• Windowills
• Counters	• Drinking Fountains	• Glazed Porcelain	• Portable Toilets	• Sports Equipment	• Wheelchairs
• Changing Tables	• Examination tables	• Linen carts	• Salad Bar Sneeze Guards	• Stainless Steel*	• Whirlpool Interiors

\* Follow directions for Toys under Directions for Use - rinsing application required. † Non-Medical or Fixture.

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) AND HEPATITIS B VIRUS ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS.** In health care settings or other settings in which there is an expected likelihood of soiling of inanimate surface/objects with blood or body fluids, and in which the surface/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 associated with AIDS and Hepatitis B Virus.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HEPATITIS B VIRUS ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:**

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and Hepatitis B Virus. For all other organisms, surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

**Intended for use only in:**

• Ambulances	• Clinics	• Hotel/Motel	• Nurseries	• Physicians Offices	• Schools
• Blood Banks	• Colleges	• Health Clubs	• Laboratories	• Nursing Homes	• Restaurants
• Cafeterias	• Day Care Centers	• Hospitals	• Military Installations	• Pharmacies	• Restrooms
				• Veterinary Offices	• Waiting Rooms

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. **This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.** Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 2 to 3 seconds until covered with mist.

- **To Disinfect:** Let stand for 10 minutes then allow to air dry.
- **To Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Sanitize:** Let stand for 30 seconds then allow to air dry.
- **To Deodorize:** Spray on surfaces as needed.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry. Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

**PRECAUTIONARY STATEMENTS:** Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**FIRST AID**

**If in eyes:** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.

**CONTACT NUMBER**

Questions? Comments or in case of an emergency, call toll free 1-800-228-4722.

Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.

**PHYSICAL HAZARDS:** FLAMMABLE. Contents under pressure. Keep away from heat, sparks and open flame.

Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

**DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

Questions? (1-800-228-4722)

For ingredient and other information, [www.rbnainfo.com](http://www.rbnainfo.com)

Distributed by: Reckitt Benckiser

Parsippany, NJ 07054-0224

Made in U.S.A. © 2012 BB

EPA Reg. No.: 777-99-675

EPA Est. No.: 777-NJ-2

(See Bottom for Lot/Date code)

0396302

Reckitt Benckiser



Federal regulations prohibit CFC propellants in aerosols.



PLEASE RECYCLE WHEN EMPTY



**KILLS COLD & FLU VIRUSES\***

**Professional Lysol®**

BRAND III

**KILLS 99.9% OF VIRUSES & BACTERIA\*\***

**Disinfectant Spray**

Eliminates odors

**Country Scent®**



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. **9833.249**

NET WT.:  
19 OZ.  
(1 LB. 3 OZ.)  
538 g

**ACTIVE INGREDIENTS:**

Alkyl (50% C14, 40% C12, 10% C16)  
dimethyl benzyl ammonium saccharinate...0.10%  
Ethanol.....58.00%  
**OTHER INGREDIENTS:**.....41.90%  
**TOTAL:**.....100.00%

KEEP OUT OF REACH OF CHILDREN  
**CAUTION:** SEE BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS

This product meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants. GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL\*

Kills the source of mold allergens.††

This product eliminates odors at their source. This product deodorizes by killing bacteria that cause odors. Ordinary, non-germicidal sprays can't do this.

**Kills in 30 seconds:**

• Avian Influenza A Virus • Campylobacter jejuni • Coxsackie Type B3 Virus • Cytomegalovirus • Hantavirus • Hepatitis B Virus • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) (AIDS Virus) • Influenza A Virus\* • Influenza B Virus • Respiratory Syncytial Virus • Rhinovirus Type 39† • Rotavirus WA • Vaccinia Virus

**Kills in 5 minutes:**

• Enterobacter aerogenes • Vancomycin Resistant Enterococcus faecalis (VRE) • Escherichia coli (O157:H7) • Listeria monocytogenes • Pseudomonas aeruginosa • Salmonella enterica • Staphylococcus aureus • Trichophyton mentagrophytes (Athlete's Foot Fungus)

**\*\*Kills in 10 minutes:**

**BACTERIA:**  
 • Acinetobacter calcoaceticus • Klebsiella pneumoniae • Proteus mirabilis • Shigella dysenteriae  
 • Burkholderia cepacia • Carbapenem Resistant • Proteus vulgaris • Staphylococcus epidermidis  
 • Corynebacterium diphtheriae • Methicillin Resistant • Pseudomonas putida • Streptococcus pyogenes  
 • Enterococcus faecalis • Staphylococcus aureus (MRSA) • Salmonella enterica serovar enteritidis • Streptococcus salivarius  
 • Klebsiella pneumoniae • Mycobacterium bovis BOG (Quant tuberculosis) • Salmonella enterica serovar Paratyphi B  
 • Klebsiella pneumoniae - NDM-1 positive • Neisseria elongata • Salmonella enterica serovar typhi  
 • Serratia marcescens

**FUNGI:**

• Aspergillus niger (mold & mildew)†† • Candida albicans • Fusarium solani • Penicillium chrysogenum††

**VIRUSES:**

• Adenovirus Type 2 • Echovirus Type 12 • Hepatitis A Virus • Poliovirus Type 1 • Feline calicivirus (Norovirus)

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces • Door handles • Floors • Patient Chairs • Shower Areas • Telephones  
 • Bed Frames • Dressing carts • Faucets • Plastic Toys\* • Sinks • Windowills  
 • Counters • Drinking Fountains • Glazed Porcelain • Portable Toilets • Sports Equipment • Wheelchairs  
 • Changing Tables • Examination tables • Linen carts • Salad Bar Sneeze Guards • Stainless Steel\* • Whirlpool Interiors

\* Follow directions For Toys under Directions for Use - rinsing application required. †† Non-Medical or Fixture

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) AND HEPATITIS B VIRUS ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HEPATITIS B VIRUS ON SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS:**

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV-1 and Hepatitis B Virus. For all other organisms, surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

Intended for use only in:

• Ambulances • Clinics • Dental Offices • Hotel/Motel • Nurseries • Physicians Offices • Schools  
 • Blood Banks • Colleges • Health Clubs • Laboratories • Nursing Homes • Restaurants • Veterinary Offices  
 • Cafeterias • Day Care Centers • Hospitals • Military Installations • Pharmacies • Restrooms • Waiting Rooms

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. **This product is not for use on critical & semi-critical medical devices or medical equipment surfaces.** Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 2 to 3 seconds until covered with mist.

- **To Disinfect:** Let stand for 10 minutes then allow to air dry.
- **To Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Sanitize:** Let stand for 30 seconds then allow to air dry.
- **To Deodorize:** Spray on surfaces as needed.
- **To Control and Prevent Mold & Mildew:** Apply to pre-cleaned surface. Allow to remain wet for 3 minutes. Let air dry.

Repeat applications in weekly intervals or when mold and mildew growth appears.

Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

**PRECAUTIONARY STATEMENTS:** Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
<b>If in eyes</b>	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
CONTACT NUMBER	
Questions? Comments or In case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.	

**PHYSICAL HAZARDS: FLAMMABLE.** Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

**DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container. To avoid waste, use all material in this container according to label directions. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

**Questions? (1-800-228-4722)**

For ingredient and other information, [www.rbnainfo.com](http://www.rbnainfo.com)

Distributed by: Reckitt Benckiser  
 Parsippany, NJ 07054-0224  
 Made in U.S.A. © 2012 RB

EPA Reg. No.: 777-99-675  
 EPA Est. No.: 777-NJ-2  
 (See Bottom for Lot/Date code)  
 0366479




Federal regulations prohibit CFC propellants in aerosols.



PLEASE RECYCLE WHEN EMPTY



**KILLS COLD & FLU VIRUSES\***



**KILLS 99.9% OF VIRUSES & BACTERIA\*\***

**Disinfectant Spray**

**Eliminates odors**

**Crisp Linen® scent**



STATE OF HAWAII  
 Department of Agriculture

**ACCEPTED**

LICENSE NO. **9833.249**

NET WT.:  
 19 OZ.  
 (1 LB. 3 OZ.)  
 538 g

**ACTIVE INGREDIENTS:**  
 Alkyl (50% C14, 40% C12, 10% C16)  
 dimethyl benzyl ammonium saccharinate...0.10%  
 Ethanol.....58.00%  
**OTHER INGREDIENTS:**.....41.90%  
**TOTAL:**.....100.00%

KEEP OUT OF REACH OF CHILDREN  
**CAUTION:** SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

This product meets AOAC Germicidal Spray Product Test standards for hospital aerosol disinfectants.  
GERMICIDAL, FUNGICIDAL, TUBERCULOCIDAL, VIRUCIDAL  
Kills the source of mold allergens.††

**Kills in 30 seconds:**

• Avian Influenza A Virus • Campylobacter jejuni • Coxsackie Type B3 Virus • Dytomegalovirus • Hantavirus • Hepatitis B Virus • Herpes Simplex Virus Type 1 & 2 • Human Immunodeficiency Virus Type 1 (HIV-1) [AIDS Virus] • Influenza A Virus • Influenza B Virus • Respiratory Syncytial Virus • Rhinovirus Type 39 • Rotavirus Wa • Vaccinia Virus

**Kills in 5 minutes:**

• Enterobacter aerogenes • Enterococcus faecalis (VRE) • Escherichia coli (O157:H7) • Listeria monocytogenes • Pseudomonas aeruginosa • Salmonella enterica • Staphylococcus aureus • Trichophyton mentagrophytes (Athlete's Foot Fungus)

**\*\* Kills in 10 minutes:**

**BACTERIA:**

• Acinetobacter calcoaceticus	• Mycobacterium bovis BCG	• Pseudomonas putida	• Staphylococcus epidermidis
• Burkholderia cepacia	• (Quant. tuberculose)	• Salmonella enterica serovar enteritidis	• Streptococcus pyogenes
• Corynebacterium diphtheriae	• Methicillin Resistant	• Salmonella enterica serovar Paratyphi B	• Streptococcus salivarius
• Enterococcus faecalis	• Staphylococcus aureus (MRSA)	• Salmonella enterica serovar typhi	
• Klebsiella pneumoniae	• Neisseria elongata	• Serratia marcescens	
• Proteus vulgaris	• Proteus mirabilis	• Shigella dysenteriae	

**FUNGI:**

• Aspergillus niger (mold & mildew)†† • Candida albicans • Fusarium solani • Penicillium chrysogenum††

**VIRUSES:**

• Adenovirus Type 2 • Echovirus Type 12 • Hepatitis A Virus • Poliovirus Type 1 • Feline calicivirus (Norovirus)

This product disinfects and sanitizes the following surfaces:

• Bathroom surfaces	• Door handles	• Floors	• Patient Chairs	• Shower Areas	• Telephones
• Bed Frames	• Dressing carts	• Faucets	• Plastic Toys*	• Sinks	• Windowills
• Counters	• Drinking Fountains	• Glazed Porcelain	• Portable Toilets	• Sports Equipment	• Wheelchairs
• Changing Tables	• Examination tables	• Linen carts	• Salad Bar Sheeze Guards	• Stainless Steel*	• Whirlpool Interiors

\* Follow directions For Toys under Directions for Use - rinsing application required. \*\* Non-Medical or Fixture

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) [AIDS VIRUS] AND HEPATITIS B VIRUS ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS) and Hepatitis B Virus.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HEPATITIS B VIRUS ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:**

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings.
- **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
- **CONTACT TIME:** Allow surface/object to remain wet for 30 seconds for HIV and Hepatitis B Virus. For all other organisms, surfaces must remain wet for 10 minutes.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

**Intended for use only in:**

• Ambulances	• Clinics	• Dental Offices	• Hotel/Motel	• Nurseries	• Physicians Offices	• Schools
• Blood Banks	• Colleges	• Health Clubs	• Laboratories	• Nursing Homes	• Restaurants	• Veterinary Offices
• Cafeterias	• Day Care Centers	• Hospitals	• Military Installations	• Pharmacies	• Restrooms	• Waiting Rooms

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using the product. This product is not for use on critical & semi-critical medical devices or medical equipment surfaces. Pre-clean surfaces prior to use. Hold can upright 6" to 8" from surface. Spray 2 to 3 seconds until covered with mist.

- **To Disinfect:** Let stand for 10 minutes then allow to air dry.
- **Rinse toys and food contact surfaces with potable water after use.**
- **For Tuberculocidal activity,** this product is effective in 10 minutes at 20°C (68°F) on surfaces that have been thoroughly cleaned prior to application.
- **To Sanitize:** Let stand for 30 seconds then allow to air dry.
- **To Deodorize:** Spray on surfaces as needed.
- **To Control and Prevent Mold & Mildew and their Odors:** Apply to pre-cleaned surface. Allow to remain wet for 10 minutes. Let air dry.

Repeat applications in weekly intervals or when mold and mildew growth appears.  
Do not use on polished wood, painted surfaces, leather, rayon fabrics, or acrylic plastics.

**PRECAUTIONARY STATEMENTS:** Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Do not spray in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID	
<b>If in eyes</b>	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
CONTACT NUMBER	
Questions? Comments or in case of an emergency, call toll free 1-800-228-4722. Have the product container or label with you when calling a Poison Control Center, or doctor, or going for treatment.	

**PHYSICAL HAZARDS:** FLAMMABLE. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage and disposal.  
**STORAGE:** Store in original container in areas inaccessible to children. Keep securely closed.

**DISPOSAL:** Do Not Puncture or Incinerate! Non-refillable container. Do not reuse or refill this container.  
If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

This can is made from an average of 25% recycled steel (10% post-consumer).

**Questions? Call 1-800-228-4722**

For ingredient and other information,

[www.rbnainfo.com](http://www.rbnainfo.com)

**IMPORTANTE: Para instrucciones de uso y primeros auxilios en español por favor llame al 1-877-843-3817.**

Distributed by: Reckitt Benckiser Inc.  
Parsippany, NJ 07054-0224  
Made in U.S.A. © RBL 2009  
EPA Reg. No.: 777-99-675  
EPA Est. No.: 054487-GA-001, 777-NJ-2  
(See Bottom for Lot/Date code)  
0248135




Federal regulations prohibit CFC propellants in aerosols.



PLEASE RECYCLE WHEN EMPTY



**KILLS COLD & FLU VIRUSES\***  
**KILLS PANDEMIC 2009 H1N1 INFLUENZA A VIRUS**



**Disinfectant Spray**

*Desinfectante en Aerosol*

**Eliminates odors**

**Crisp Linen®**  
*scent*



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. **9833.249**

NET WT.:  
19 OZ.  
(1 LB. 3 OZ.)  
538 g

**ACTIVE INGREDIENTS:**

Alkyl (50% C14, 40% C12, 10% C16)	
dimethyl benzyl ammonium saccharinate...	0.10%
Ethanol.....	58.00%
OTHER INGREDIENTS.....	41.90%
<b>TOTAL:</b>	<b>100.00%</b>

KEEP OUT OF REACH OF CHILDREN  
**CAUTION:** SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
MANTENGASE FUERA DEL ALCANCE DE LOS NIÑOS  
**PRECAUCIÓN**







