Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

9903.30
Disc.

Preen.

LAWN

BROADLEAF WEED CONTROL

COVERS 5000 SQ. FT.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2008-2010 LIC. NO.

LICENSED
HAWAII
DEPT. OF AGRICULTURE
Period 2007-2009 No. 9903.30

COVERS 5000 SQ. FT.

Preen®

LAWN

BROADLEAF WEED CONTROL

- Controls dandelions and 200 other broadleaf weeds
- Will not burn your lawn
- Apply anytime weeds are growing - spring, summer or fall

COVERS
5,000 SQ. FT.
Net Wt. 10 lbs. (4.54 kg)
Does not contain fertilizer

Keep out of reach of children
CAUTION
See back panel for Precautionary Statements

Good Housekeeping Promise

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid	1.37%
(+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.31%
Dicamba: 3,6-dichloro-o-anisic acid	0.13%
INERT INGREDIENTS	98.19%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
2.1 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton
0.2 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
2.6 lbs. 3,6-dichloro-o-anisic acid equivalent per ton

Same as Specific by AOAC Methods
Contains the single isomer form of Mecoprop-p

Preen.

WEED CONTROL

- Controls dandelions and 200 other broadleaf weeds
- Will not burn your lawn
- Apply anytime weeds are growing - spring, summer or fall

COVERS
5,000 SQ. FT.
Net Wt. 10 lbs. (4.54 kg)
Does not contain fertilizer

Keep out of reach of children
CAUTION
See back panel for Precautionary Statements.

Good Housekeeping Promises

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid	1.37%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid	0.31%
Dicamba: 3,6-dichloro-o-anisic acid	0.13%
INERT INGREDIENTS:	98.19%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
6.2 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.
2.6 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods
Contains the single isomer form of Mecoprop-p.

Preen®

LAWN BROADLEAF WEED CONTROL

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

Use Precautions

Do not allow the material to drift onto flowers, vegetables or ornamental shrubbery. Apply when the air is calm. Do not apply to newly seeded lawns. Do not apply in areas underlaid by roots of desirable trees and shrubs. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

Instructions

Preen Lawn Broadleaf Weed Control controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

Established turf of warm season grasses — includes bermudagrass, centipedegrass, and zoysiagrass.

Established turf of cool season grasses — includes Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues.

Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing desirable clovers.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Spreader Settings

Weeds

Annual yellow sweetclover
Aster
Austrian fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, field
Bird vetch
Bitter wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Black mustard
Blackseed plantain
Blessed thistle
Bloodflower milkweed
Blue lettuce
Blue vervain
Bracted plantain
Bristly oxtongue
Broadleaf dock
Broadleaf plantain
Broomweed
Buckhorn
Buckhorn plantain
Bulbous buttercup
Bull thistle
Bullnettle
Burclover
Burdock
Burning nettle
Burweed
Cocklebur
Common mullein
Creeping jenny
Cudweed
Curly dock
Daisy: English, oxeye
Daisy fleabane
Dandelion
Dichondra
Doglennel
Dollarweed
False dandelion
Falseflax
False sunflower
Fiddleneck
Florida pusley
Frenchweed
Galinsoga
Goldenrod
Ground ivy
Gumweed
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Heath aster
Hedge bindweed
Hedge mustard
Henbit
Hoary cress
Hoary plantain
Hoary vervain
Horsenettle
Jimsonweed
Knawel
Orange hawkweed
Oriental cocklebur
Oxalis
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coreopsis
Plantain
Poison ivy
Poison oak
Pokeweed
Poorjoe
Prairie sunflower
Prickly lettuce
Prickly sida
Prostrate knotweed
Prostrate pigweed
Prostrate spurge
Prostrate vervain
Puncturevine
Purslane, common
Ragweed
Red sorrel
Redroot pigweed
Redstem filaree
Rough cinquefoil
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimp
Spotted spurge
Spurweed
Stinging nettle
Strawberry, India mock
Tall nettle
Tall vervain
Tansy ragwort
Tansy mustard
Tanweed
Thistle
Trailing crownvetch
Tumble mustard
Tumble pigweed
Velvetleaf
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify
White mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild marigold
Wild mustard
Wild onion
Wild parsnip
Wild radish

The spreader settings below are approximate. Calibrate your spreader on 100 sq. ft. and adjust to apply at the rate of 1 lb. of product per 500 sq. ft. (2 lbs. of material per 1,000 sq. ft.). Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.



Apply in a single pass at a rate of 2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.

Broadcast Spreaders	Setting
Agri-Fab® Rotary	5
Cyclone® Rotary	4
Earthway® Rotary	16
Greenview® Jet-Spred® SS1	16
Lesco® Rotary (dial A-S)/(number dial)	G /15
Red Devil® Rotary (1-20)	5
Republic® Rotary	2
Scotts® EasyGreen®	26
Scotts® EdgeGuard®	5
Scotts® SpeedyGreen® 2000, 3000	5
Sears® Craftsman Rotary	5
Sta-Green® Rotary (1-10)	3
True Temper® Rotary	4
Vigoro® Rotary (1-10)	3



Apply in a single pass at a rate of 2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.

Drop Spreaders	Setting
Agri-Fab® Drop	4
Greenview® Drop-Spred® XP1	16
Red Devil® Drop (1-5)	4
Republic® Drop	5½
Scotts® Drop AccuGreen (2-18)	5½
Scotts® Drop PF1 (2-18)	7½
Sta-Green® Drop (1-5)	4

If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

Buttercup
Buttonweed
Canada thistle
Carolina geranium
Carpetweed
Catchweed bedstraw
Catnip
Catsear
Chickweed: common, mouseear
Chicory
Cinquefoil
Clover: crimson, hop, red, strawberry, sweet, white
Cockle
Knotweed
Kochia
Lambsquarters
Lespedeza
Mallow
Matchweed
Mexicanweed
Milk vetch
Morningglory
Mouseear hawkweed
Mugwort
Musk thistle
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny amaranth
Spiny cocklebur
Spiny sowthistle
Spotted catsear
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Woodsorrel
Woolly croton
Woolly morningglory
Woolly plantain
Wormseed
Yarrow
Yellow rocket
Yellowflower
pepperweed
& other broadleaf weeds

Warranty

The Distributor warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Distributor, and Buyer assumes the risk of any such use.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.
After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

FIRST AID

IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free (888) 208-1368.

Environmental Hazards

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not contaminate domestic or irrigation waters. Do not apply when weather conditions favor drift from target area.

Storage and Disposal

Storage: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.
Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

EPA Reg. No. 2217-940-961
(L) (D)
EPA Est. 961-PA-01;961-IL-01
Superscript used is first letter of lot number.

0 88685 63696 7

PROOF OF PURCHASE

PREEN GUARANTEE

Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem, and send with proof of purchase to:

LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042

1-800-233-1067

www.preen.com

Preen® is a registered trademark of Lebanon Seaboard Corporation. 1/04