



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

9903.41



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

GREENVIEW®

CRABGRASS CONTROL PLUS LAWN FOOD

For control of crabgrass and control or suppression of many other annual grass and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf.



Enhanced Efficiency Fertilizer

GREENSMART

For a Green Lawn

Helps lawns retain water to
protect against heat & drought

Speeds root growth to
improve nutrient absorption

Feeds your lawn up to 12 weeks

Apply in Spring



For more information, visit www.greenviewfertilizer.com

PREVENTS CRABGRASS ALL SEASON

In the state of New York, do not exceed 3.0 lb./1000 sq. ft. per year (equivalent to 0.25 pounds of active ingredient per acre).

NET WEIGHT 13.5 lbs. (6.12 kg)
Covers 5,000 Sq. Ft.

ACTIVE INGREDIENT: 0.19%
Dithiopyr.....
OTHER INGREDIENTS: 99.81%
TOTAL 100.00%

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See back panel for First Aid and
Additional Precautionary Statements



GREENVIEW® CRABGRASS CONTROL

PLUS LAWN FOOD

PREVENTS CRABGRASS ALL SEASON

ertilizer.com

In the state of New York, do not exceed 3.0 lb./1000 sq. ft. per year
(equivalent to 0.25 pounds of active ingredient per acre).

lbs. (6.12 kg)
Ft.

ACTIVE INGREDIENT:
Dithiopyr..... 0.19%
OTHER INGREDIENTS:..... 99.81%
TOTAL 100.00%

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See back panel for First Aid and
Additional Precautionary Statement

NOTICE: Read the entire label. Use only according to label directions. Before buying this product, read the Warranty Section elsewhere on this label. If terms are not acceptable, return the unopened container at once.

**CARE & HANDLING
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Treatment Information call (888)208-1368.	

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies and drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
For Outdoor Residential Use.
Read all Directions for Use carefully before applying.

How This Product Works

GreenView Crabgrass Control plus Lawn Food can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months of crabgrass control.

GreenView Crabgrass Control plus Lawn Food provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

<u>Cool-Season Grasses</u>	<u>Warm-Season Grasses</u>
Bentgrass, creeping	Bahiagrass St. Augustinegrass
Bluegrass, Kentucky	Bermudagrass Zoysiagrass
Fescue, fine	Buffalograss
Fescue, tall	Carpentgrass
Ryegrass, perennial	Centipedegrass

Use Precautions

GreenView Crabgrass Control plus Lawn Food must be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, use the initial application of this product only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. Exception: In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

Do not use on turf grown for agricultural or commercial purposes (turf farms), or for grass seed production.

- Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.
- For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.
- Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 6.0 lb./1000 sq. ft. (0.5 lb. active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.
- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.

Do not allow people or pets to enter treated area until grass has returned.

Weeds Controlled or Suppressed

GreenView Crabgrass Control plus Lawn Food will provide control of crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior their germination.

Grasses-	brome	crowfootgrass †	foxtail, yellow	perennial)
barley	crabgrass, large	dallisgrass	kikuyugrass †	sandbur
barnyardgrass	crabgrass, smooth	goosegrass	oats, wild	smutgrass
bluegrass, annual	crabgrass, Southern	foxtail, green	ryegrass (annual &	
Broadleaf Weeds-	henbit	mustard	purslane, common	spurge, prostrate
bittercress †	knotweed, prostrate	oxalis, buttercup	rocket, London	spurge, spotted
carpetweed	lespedeza, common †	pineappleweed	shepherdspurse	woodsorrel, creeping
chickweed †	marestail	pigweed, redroot	speedwell, corn †	woodsorrel, yellow
geranium, Carolina †	medic, black	parsley-piert †	spurge, garden	

† Suppression only

How to Apply

To optimize product performance apply this product within a few days after mowing and delay mowing again for a few days after the application. Most effective weed control will be obtained when GreenView Crabgrass Control plus Lawn Food is activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.


SPREADER SETTINGS

Broadcast Spreader	Setting				Drop Spreader	Setting	
	2.7 Lbs./1,000 Sq. Ft.					2.7 Lbs./1,000 Sq. Ft.	
Agri-Fab® Rotary	4	Scotts Edgeguard	4½	Agri-Fab Drop	4		
Cyclone Rotary	4½	Scotts Speedy Green 1000	4½	Greenview® Drop-Spreader® XP1	15		
Earthway Rotary	14	Scotts Speedy Green 2000	4½	Red Devil Drop (1-5)	3		
Greenview Jet-Spreader SS1	14	Scotts Speedy Green 3000	4½	Republic Drop	5		
Lesco (letter/numeric)	G/15	Sears Craftsman Rotary	4	Scotts AccuGreen	5		
Red Devil Rotary (1-20)	3½	Sta-Green Rotary (1-10)	3½	Scotts Drop PF1 (2-18)	4½		
Republic Rotary	4	True-Temper Rotary	3½	Sta-Green Drop (1-5)	3½		
Scotts EasyGreen	22	Vigoro Rotary (1-10)	3½				


These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Spreader Directions:


Broadcast Spreader: Apply in a single pass by starting on the outside border and apply in a circular path working toward the center.



Drop Spreader: First make two passes along the top of the area you'll be covering. Then make two passes along the bottom area. Then move back and forth along the long part of the lawn. On each pass, as you reach the strips at the top and bottom of the area to be covered, shut off the spreader. It's also important to shut off the spreader when you are stopped.



Determining Your Lawn's Square Footage



To determine how much of this product to apply, you must determine the square footage of your lawn. An easy way to do this is to walk the length of your lawn, figuring that one pace equals about 3 feet. Do the same with the width of the lawn. Then multiply the length by the width to arrive at the total. Make sure you subtract the square footage of your home and driveway when calculating the total square footage of your lawn. If your lawn has an irregular shape, try dividing it up into a number of square areas. Calculate the square footage for each area, then add the figures together to reach your total lawn size.

Spreader Tips


The two main types of spreaders are the Drop Spreader, which drops the material straight down the width of the spreader as you push, and the Broadcast Spreader (also known as a rotary), which throws the material from the spreader. The above spreader settings are approximate. To check your spreader: 1 lb. of material covers approximately 370 sq. ft. (2.7 lbs. of material per 1,000 sq. ft.)

Note: if spreader model does not appear above please call toll free 800-233-1067, 8 am - 5 pm ET, Monday - Friday for additional model settings.

How Much To Apply

GreenView Crabgrass Control plus Lawn Food will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied.

Refer to the map below to determine the region of the U.S. that includes your location. Use the table below to determine the amount of GreenView Crabgrass Control plus Lawn Food to apply in your location based on the desired length of weed control or need for postemergence crabgrass control. Use "regions" within the United States:



Product Use Rates

Amount of GreenView Crabgrass Control plus Lawn Food to Apply (Pounds of product to apply per 1,000 square feet of turfgrass [†])			
Region of U.S.	----- Length of Crabgrass Control -----		
	2 - 3 months	3 - 4 months	Early Postemergence Control
North	1.5 lb./1000 sq. ft. (0.128 lb. ai/acre)	2.1 lb./1000 sq. ft. (0.174 lb. ai/acre)	2.7 lb./1000 sq. ft. (0.22 lb. ai/acre)
South	2.3 lb./1000 sq. ft. (0.192 lb. ai/acre)	2.7 lb./1000 sq. ft. (0.22 lb. ai/acre)	2.7 lb./1000 sq. ft. (0.22 lb. ai/acre)

[†] 1,000 square feet equals an area 100 feet X 10 feet or square area of about 32 feet on a side.

Maximum Application Rates:

- Do not apply more than 6.0 lb. of this product per 1,000 sq. ft. per application and no more than 18.0 lb. of this product/1000 sq. ft. per year (equivalent to 0.5 and 1.5 lb. of active ingredient per acre, respectively).
- In the state of New York, do not exceed 3.0 lb./1000 sq. ft. per year (equivalent to 0.25 pounds of active ingredient per acre).

WHEN TO APPLY

Spring Application

Control of Crabgrass: GreenView Crabgrass Control plus Lawn Food provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

Control or Suppression of Other Grass and Broadleaf Weeds: For control or suppression of other grass and broadleaf weeds, this product must be applied as a preventative treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

Fall Application

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, do not reseed, overseed or sprig turfgrasses within 12 weeks after application.

spring turgrasses within 12 weeks after application.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store this product only in its original container in a cool, dry, locked place out of the reach of children and animals.

Pesticide Disposal and Container Handling: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Lebanon Seaboard Corporation. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, LEBANON SEABOARD CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Lebanon Seaboard Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Lebanon Seaboard Corporation DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT LEBANON SEABOARD CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

GUARANTEED ANALYSIS 26-0-4

Total Nitrogen (N)	26%
2.4% Ammoniacal Nitrogen	
1.5% Water Insoluble Nitrogen*	
19.6% Urea Nitrogen	
2.5% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	4%
Sulfur (S)	2.6%
2.6% Combined Sulfur (S)	
Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.	
Chlorine (Cl) not more than	4.0%
* 4.0% Slowly Available Nitrogen from Methylene Ureas.	F699

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-lebsea.com>.



Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem and simply send proof of purchase to Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.

- For more information on our complete line of GreenView® Lawn and Garden Products, visit our website at www.greenviewfertilizer.com
- GreenView® is a registered trademark of Lebanon Seaboard Corporation.

Manufactured by:
Lebanon Seaboard Corporation
1600 East Cumberland Street
Lebanon, PA 17042
Made in the USA

21-31159

EPA Reg. No. 961-417

(L) (D) (BE)
EPA Est. Nos. 961-PA-01, 961-IL-01, 59144-GA-1
Superscript is first letter(s) of lot number.