

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Goal[®] 2XL

EPA Reg. No. 62719-424

EPA SLN No. HI-020005

For Preemergence and Postemergence Weed Control in Eucalyptus Plantings

(For Distribution and Use Only in the State of Hawaii)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read this SLN labeling and the label affixed to the container for Goal[®] 2XL herbicide before applying. All applicable use directions, precautions and restrictions on this SLN labeling and the label affixed to the product container must be followed.

Directions for Use

Note: Participation by the Applicator (i.e., supervisor or person responsible for application) in Dow AgroSciences' Stewardship Program for the application of Goal 2XL in Eucalyptus plantings is required prior to use.

General Information: Goal[®] 2XL herbicide may be applied for postemergence and preemergence control of certain broadleaf weeds in permanently established eucalyptus (such as *Eucalyptus viminalis*, *E. pulverulenta*, and *E. camaldulensis*) plantings. If used on species not specifically listed, or if unsure of the species tolerance to Goal 2XL, test for crop safety on a small number of plants before proceeding.

Application Rates: Apply Goal 2XL at 4 to 8 pints (1.0 to 2.0 lb active) per broadcast acre for preemergence and postemergence weed control. A maximum of 16 pints of Goal 2XL (4.0 lb active) per acre per season may be applied as a result of multiple applications. For postemergence applications, the addition of an 80% active non-ionic surfactant cleared for use on crops at the rate of 1 quart per 100 gallons of spray mix will improve spray coverage and wetting of weeds.


Weeds Controlled: When Goal 2XL is applied preemergence or postemergence to weed seedlings (not exceeding 6-leaf stage) at recommended application rates, the following broadleaf weeds are controlled:

Weeds Controlled Postemergence

Cheeseweed (Malva)	Minerslettuce	Filaree, Whitestem [†]	Shepherdspurse
Groundsel, Common	Shepherdspurse	Pigweed, Redroot	Sowthistle, Annual
Purslane, Common ^{††}	Filaree, Redstem [†]	Spurge, Garden ^{††}	Spurge, Garden
Fiddleneck, Coast	Nettle, Burning	Redmaids	Spurge, Prostrate
Henbit Redmaids	Sowthistle, Annual	Rocket, London	Spurge, Spotted
Filaree, Broadleaf [†]			

[†] Goal 2XL at the 8-pint rate (2.0 lb active) per acre will provide control of filaree up to the 6-leaf stage.

^{††} Seedlings beyond the 4-leaf stage may result in partial control.



Weeds Controlled Preemergence

Ageratum	Knotweed, Prostrate	Filaree, Broadleaf
Burclover	Crotalaria	Pigweed, Redroot
Groundsel, Common	Lambsquarters, Common	Filaree, Redstem
Buttonweed Henbit	Fiddleneck, Coast	Purslane, Common
Cheeseweed (Malva)	Lettuce, Prickly	

Timing and Method of Application: In new plantings, Goal 2XL should be applied immediately before or after transplanting of dormant eucalyptus seedlings. In established plantings, Goal 2XL may be applied postemergence (over the top or post-directed to the base of the eucalyptus trees). It is recommended that application be made prior to bud break or as a post directed application to avoid/reduce possible phytotoxicity to the eucalyptus foliage. It is recognized that Goal 2XL can be applied postemergence (over-the-top) to non-dormant eucalyptus plantings, however, growers should understand that foliage receiving direct or indirect (drift) sprays may exhibit unacceptable crop injury.

Apply Goal 2XL at 20 to 40 psi in a minimum of 20 gallons of spray volume per acre. Increase the spray volume as weed height and density increase. Use a low-pressure sprayer equipped with flat fan nozzles. Check calibration of spray equipment before use.

Refer to main product label for Mixing Directions.

Crop-Specific Use Restrictions:

The following use restrictions must be observed when Goal 2XL is used as recommended on this label:

- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EP A registered label.
- Do not apply more than 8 pints (2.0 lb active) of Goal 2XL per broadcast acre per application or more than 16 pints (4.0 lb active) per broadcast acre per year.
- Make applications of Goal 2XL only to healthy trees.
- Do not allow grazing or harvest forage from areas treated with Goal 2XL.
- Goal 2XL is phytotoxic to plant foliage. Do not apply when weather conditions favor drift (i.e. when wind speed exceeds 10 mph). Applicator must have means to accurately measure wind speed during application. Avoid drift to all non-target areas.
- Do not treat ditch banks or waterways with Goal 2XL.
- Aerial application is prohibited.
- **Chemigation:** For uses described in this supplemental labeling, do not apply this product through any type of irrigation system.

24(c) Registrant: Dow AgroSciences • 9330 Zionsville Road • Indianapolis, IN 46268

This label is valid until January 27, 2012 or until otherwise amended, withdrawn, cancelled or suspended.

Issue Date: January 28, 2007

Expiration Date: 01-27-2012

®Trademark of Dow AgroSciences LLC

R-843-013
Accepted: 01-09-07
Replaces D06-843-064

