

Goal[®] 2XL

EPA Reg. No. 62719-424

EPA SLN No. HI-020010

For Preemergence and Postemergence Weed Control in Banana Plantings**(For Distribution and Use Only in the State of Hawaii)****ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read this SLN labeling and the label affixed to the container for Goal[®] 2XL herbicide before applying. All applicable use directions, precautions and restrictions on this SLN labeling and the label affixed to the product container must be followed.

Directions for Use

Apply Goal[®] 2XL herbicide as a directed application for preemergence and postemergence broadleaf weed control in permanently established bearing and nonbearing banana plantings.

Method of Application: Banana plants must be healthy and of sufficient height to allow for application of directed sprays (greater than 2 feet tall). Sprays should be directed to the orchard floor beneath banana plants. Prevent direct spray contact with banana plants, including foliage, as injury may result.

Use a low-pressure sprayer equipped with a breakaway boom and flat fan or off-center (OC) nozzles. Apply Goal 2XL in a minimum spray volume of 20 gallons per acre. Minimize potential for spray drift to banana plants by using nozzles that produce coarse sprays and minimum spray pressure recommended by the manufacturer. Use higher spray volumes to ensure adequate coverage if emerged weed density is high or if heavy crop residue (trash) is present. Check calibration of spray equipment before use.

Refer to main product label for Mixing Directions and instructions for band application.

Postemergence Application: Rates and Weeds Controlled: Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. Use a higher rate in rate range for heavy weed infestations, weeds in advanced stages of growth or where a longer period of residual weed control is desired. Applications to weeds beyond the four-leaf stage may result in partial control. The addition of an 80% non-ionic surfactant cleared for use on crops at the rate of 1 quart per 100 gallons of spray mix will improve spray coverage and wetting of weeds for postemergence applications.

Purslane, common
Spurge, garden

Tank Mixing: For postemergence control of certain grass and broadleaf weeds, a tank mixture of either paraquat (Gramoxone Extra Herbicide) or glyphosate with Goal 2XL may be applied as a directed spray to seedling weeds. Follow all applicable use directions, application rates, use precautions and restrictions on respective tank mix product labels. In interpreting each label for the tank mixture, the most restrictive limitations must apply. Do not allow sprays containing paraquat or glyphosate to contact banana stems or foliage as severe injury or crop destruction may occur.

ACCEPTED

February 7, 2012

**Under Hawaii Pesticides Law
as Supplement to Product No.
9786.189**

When Goal 2XL is tank mixed with paraquat or glyphosate and applied postemergence the following additional weeds are controlled:

Ageratum
Buttonweed
Crotalaria

Preemergence Application: Rates and Weeds Controlled: Applications for preemergence weed control are most effective when applied to final soil surfaces that are clean and weed-free. For best preemergence activity, at least one-quarter inch (1/4 inch) of water should be applied within 3 to 4 weeks after application if no rainfall occurs. Cultural practices that disturb or redistribute surface soil such as such as cutting water furrows or cultivation that mixes untreated and treated soil will reduce the herbicidal effectiveness. Best results are obtained when established berms or soil surfaces are left undisturbed during the time period for which weed control is desired.

Apply 2 to 8 pints (0.5 to 2.0 lb active ingredient) of Goal 2XL per broadcast acre. Use higher rates in the rate range on medium to fine texture soils, soils containing greater than 1% organic matter, or where a longer period of residual weed control is desired.

Ageratum	Purslane, common
Buttonweed	Spurge, garden
Crotalaria	

Crop-Specific Use Restrictions:

- Follow all applicable use directions, precautions, restrictions, Worker Protection Standard requirements on the EPA registered label.
- Do not apply more than 8 pints (2.0 lb active) of Goal 2XL per broadcast acre per year as a result of a single or multiple applications.
- **Preharvest Interval:** Goal 2XL may be applied up to 1 (one) day prior to harvest.
- Do not allow animals to graze or forage areas treated with Goal 2XL.
- Apply only as a directed spray using ground application equipment.
- Goal 2XL is phytotoxic to plant foliage. Do not allow sprays or drift to contact banana stems or foliage. Avoid drift to non-target areas.
- Do not apply Goal 2XL to ditch banks or waterways.
- **Chemigation:** Do not apply this product through any type of irrigation system for uses described in this supplemental labeling.

24(c) Registrant: Dow AgroSciences • 9330 Zionsville Road • Indianapolis, IN 46268

This label is valid until February 6, 2017 or until otherwise amended, withdrawn, cancelled or suspended

Issue Date: February 7, 2012

Expiration Date: February 6, 2017

®Trademark of Dow AgroSciences LLC

R843-115
Accepted:
Replaces R843-014