



**MAKHTESHIM**  
**A G of AN**  
**NORTH AMERICA**

**RESTRICTED USE PESTICIDE**

Due to acute toxicity to humans, aquatic organisms, and avian species. For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

**THIONEX 50W**  
**EPA REG. NO. 66222-62**  
**SOUTHERN GREEN STINK BUG IN**  
**MACADAMIA NUTS**  
**IN THE STATE OF HAWAII**  
**EPA SLN NO. HI-030001**

**SUPPLEMENTAL LABELING FOR DISTRIBUTION AND USE ONLY**  
**IN THE STATE OF HAWAII**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered product label.

This labeling must be in the possession of the user at the time of pesticide application.

<b>CROP</b>	<b>PEST</b>	<b>MAXIMUM RATE OF APPLICATION</b>
Macadamia Nuts	Southern Green Stink Bug	2 pounds of Thionex 50W per 100 gallons of water, apply up to 200 gallons of finished spray per acre.

Use 2 pounds of Thionex 50W per 100 gallons of water. Applications should be made as a directed spray to foliage.

**RESTRICTIONS**

- Do not use more than 4 pounds of this product (2 lbs. active ingredient endosulfan) per acre per year.
- Do not make more than 1 application per year
- Do not apply within 2 days of harvest.
- Do not apply by air.
- Do not apply this product through any type of irrigation system.
- For airblast application, the use of a closed cab is required.
- Restricted-Entry Interval (REI) of 2 days.
- A 30 foot vegetative buffer strip must be maintained between all areas treated with this product and rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds.
- This product is extremely toxic to fish and aquatic invertebrates and toxic to birds and mammals. Do not apply within 100 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds. Apply as a directed spray to foliage. If using airblast, turn off outward pointing nozzles at row ends and outer rows. Apply only when the wind speed is 3-10 mph at the application site as measured by an anemometer outside the orchard on the upwind side.

**PRECAUTIONS**

The area being treated must be vacated by unprotected persons.

**24(c) Registrant:**

Makhteshim Agan of North America, Inc.  
 4515 Falls of Neuse Rd., Suite 300  
 Raleigh, NC 27609

This labeling is valid until March 11, 2013 or until otherwise amended, withdrawn, cancelled, or suspended.

Issue Date: March 12, 2008  
 Expiration Date: March 11, 2013

