

ACCEPTED

OCT 14 2009

Under Hawaii Pesticides Law
as Supplement to Product No.
9786.246

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

GoalTender[®]

EPA Reg. No. 62719-447

EPA SLN No. HI-090003

For Preemergence and Postemergence Weed Control in Macadamia Nut Plantings (*Macadamia ternifolia* var. *integrifolia*)

(For Distribution and Use Only in the State of Hawaii)

Conditions and Risks of Use for Special Local Need

USE OF GoalTender[®] HERBICIDE (THE "PRODUCT") ON MACADAMIA NUTS (THE "CROP") FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP.

This Product is available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that, to the extent permitted by law:

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Dow AgroSciences does not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental labeling;
- (3) Dow AgroSciences does not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- (4) Dow AgroSciences disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) these Conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Dow AgroSciences, or its respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

These Conditions and Risks of Use for Special Local Need are required by Dow AgroSciences and not specified by the US EPA or the State of Hawaii.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read this SLN labeling and the label affixed to the container for GoalTender® herbicide before applying. All applicable use directions, precautions and restrictions on this SLN labeling and the label affixed to the product container must be followed.

Directions for Use

Product Information

Apply GoalTender® herbicide as a directed spray application for preemergence and postemergence broadleaf weed control in bearing and nonbearing macadamia nut plantings. Apply only to established plantings of healthy growing trees and after new foliage growth has hardened off.

Method of Application

Sprays should be directed to the orchard floor. Minimize spray contact with stems and prevent direct spray contact with foliage as injury may result.

Use a low-pressure sprayer and flat fan or off-center (OC) nozzles. Apply GoalTender in a minimum spray volume of 20 gallons per acre using a minimum spray pressure of 20 psi. Higher spray volumes may be used to ensure adequate coverage if emerged weed density is high or if heavy crop residue is present. Spray equipment should be calibrated carefully before each use.

Refer to main product label for Mixing Directions and instructions for band application.

Postemergence Application: Rates and Weeds Controlled

Apply 1 to 4 pints (0.5 to 2.0 lb active) of GoalTender per broadcast acre. On a lava soil, do not exceed 2 pints (1.0 lb active) per acre of GoalTender in a single application. The higher rate (2.0 lb active) is recommended for control of seedling weeds up to the 4-leaf stage and will provide preemergence control of susceptible weeds. Applications to weeds beyond the four-leaf stage may result in partial control. The addition of an 80% non-ionic surfactant registered for use on crops at the rate of 1 quart per 100 gallons of spray will improve spray coverage and wetting of weeds for postemergence applications.

Weeds Controlled Postemergence: common purslane; garden spurge

Tank Mixing: For postemergence control of certain grass and broadleaf weeds, apply in tank mixture with glyphosate, glufosinate, paraquat or other registered contact herbicides, as a directed spray to seedling weeds. Follow all applicable use directions, application rates, use precautions, and restrictions on the respective tank mix product labels. In interpreting each label for the tank mixture, the most restrictive limitations must apply. Do not allow these sprays to contact stems or foliage, as severe injury or crop destruction may occur.

When GoalTender is tank mixed with glyphosate, glufosinate or paraquat and applied postemergence, the following additional weeds are controlled: ageratum, buttonweed, crotalaria.

Preemergence Application (Rates and Weeds Controlled):

Apply 1 to 4 pints of GoalTender (0.5 to 2.0 lb active) per broadcast acre. Use higher rates in the rate range on medium to fine texture soils, soils containing greater than 1% organic matter, or where a longer period of residual weed control is desired. On lava soil, do not exceed 2 pints (1.0 lb. active) per acre of GoalTender in a single application.

Applications for preemergence weed control are most effective when applied to soil surfaces that are clean and weed-free. Best results are obtained when treated soil surfaces are left undisturbed for the time the weed control is desired. Cultural practices that disturb or redistribute surface soil will reduce herbicidal effectiveness.

Weeds Controlled Preemergence: ageratum, buttonweed, crotalaria, common purslane, garden spurge

Crop-Specific Use Restrictions:

- Follow all applicable use directions, precautions, restrictions, Worker Protection Standard requirements on the EPA registered label.
- Do not apply more than a total of 2.0 lbs oxyfluorfen active ingredient per acre per year. On a lava soil, do not exceed 2 pints (1.0 lb. active) per acre of GoalTender per year.
- **Preharvest Interval:** GoalTender may be applied up to 7 days before harvest. Although permitted while nuts are on the ground, treatments should be made immediately after harvest to minimize the number of nuts on the ground at the time of application.
- Do not feed or graze animals on any area treated with GoalTender.
- Apply only as a directed spray using ground application equipment.
- GoalTender is phytotoxic to plant foliage. Do not allow sprays or drift to contact foliage. Avoid drift to non-target areas.
- Do not apply GoalTender to ditch banks or waterways.
- **Chemigation:** For uses described in this supplemental labeling, do not apply this product through any type of irrigation system.

This label is valid until December 4, 2012 or until otherwise amended, withdrawn, cancelled or suspended.

Issue Date: October 14, 2009

Expiration Date: December 4, 2012

®Trademark of Dow AgroSciences LLC

R204-010
Accepted: 10/14/09
Initial printing