



DEPARTMENT OF HEALTH

**NEWS RELEASE**

**LINDA LINGLE**  
GOVERNOR

---

CHIYOME LEINAALA FUKINO M.D.  
DIRECTOR  
Phone: (808) 586-4410  
Fax: (808) 586-4444

---

For Immediate Release: February 12, 2008

08-14

**DOH CONFIRMS CASE OF MEASLES ON HAWAIIAN AIRLINES FLIGHT**

HONOLULU – The Hawai'i State Department of Health (DOH) has confirmed measles illness in an airline passenger who arrived in Honolulu from San Diego on Saturday, February 9, 2008 on Hawaiian Airlines flight #15. The passenger, a 10-month-old infant, was exposed to a confirmed case of measles in a San Diego medical office on January 25, had the onset of initial symptoms on February 7, and a rash on February 10. The illness was confirmed by the DOH State Laboratory on February 11.

Last night, the DOH began contacting all passengers at telephone numbers provided by the airline to notify them of their potential exposure. Passengers are advised to consult their physicians regarding their susceptibility to measles. A 5-month-old Hawai'i resident was exposed to the case and was administered immune serum globulin by DOH to prevent measles infection.

“So far, four additional infants, potentially exposed on the flight, have been identified and will be administered preventive treatment by DOH today,” State Epidemiologist Dr. Paul Effler said.

“Hawaiian Airlines has been extremely supportive in this effort by providing passenger contacting information quickly.”

Measles is a highly contagious viral infection characterized by fever, cough, runny nose, conjunctivitis, and a rash. It is transmitted by direct contact with droplets from a patient's respiratory tract or, less commonly, by airborne spread. Patients are contagious from 1 to 2 days before onset of symptoms (3-5 days before the rash) to 4 days after appearance of the rash. The incubation period is generally 8 to 12 days, with a range from 7-18 days after exposure.

Complications of measles include ear infection, pneumonia, and occasionally encephalitis. Death

**(more)**

occurs in 1 to 3 of every 1,000 cases reported in the United States. Fatality rates are increased in children younger than 5 years of age and immunocompromised children.

Individuals born prior to 1957 or who have two documented measles vaccinations or laboratory evidence of immunity or documentation of physician-diagnosed measles are considered immune to measles. All other passengers on board this flight should be considered potentially susceptible and exposed, and should immediately consult their physician.

**###**

For information contact:  
Janice Okubo  
Department of Health  
Phone: (808) 586-4442  
Cell: (808) 368-6006



## Hawaii State Department of Health

### Disease Investigation Branch

Oahu 586-4586, Maui 984-8213, Hilo 933-0912,

Kona 322-4877, Kauai 241-3563

---

## Measles

(Rubeola)

### What is measles?

Measles is a highly contagious rash illness caused by a virus. Although measles is commonly thought of as a childhood disease, people of any age can get it. Possible complications of measles are pneumonia, inflammation of the brain (encephalitis), ear infections, diarrhea, seizures and death. These complications are more common in children under age 5 and adults over age 20. Measles can be prevented with a vaccination.

### How do you get it?

Measles is spread through the air by droplets and by direct exposure to mucus from the nose and throat of a person infected with the measles virus. The virus can remain contagious on surfaces for up to 2 hours.

### What are the symptoms of measles?

The symptoms usually occur in two stages. The first stage begins with a high fever, runny nose, cough and conjunctivitis (pink eye). In the second stage, a red blotchy rash appears and usually lasts for 5 to 6 days. The rash commonly begins on the face and then spreads to the rest of the body.

### When do symptoms start?

The fever, runny nose, and cough usually appear 10 days after exposure to the virus, but the onset can range from 7 to 18 days. The rash follows 3 to 7 days after the beginning of the first symptoms.

### For how long is a person with measles contagious?

A person is contagious from just before the first symptoms appear to 4 days after the start of rash.

### What is the treatment for measles?

There is no specific treatment for measles. Care of patients with measles consists mainly of providing plenty of water and food, bed rest, and acetaminophen (Tylenol) for fever control.

### Is there a vaccine for measles?

Yes. All children should receive 2 doses of the measles, mumps and rubella (MMR) vaccine. The first dose should be given at 12-15 months of age. A second dose is recommended at 4 to 6 years of age. Measles vaccination is required for all children before enrollment in any school in Hawaii (preschool through college). Unvaccinated adults born after 1956 should receive at least one dose of measles vaccine, unless they have had a blood test showing they are immune to measles. Adults at increased risk of getting measles (college students, international travelers, and health care workers) should receive two doses, at least 4 weeks apart.

### Should a person with measles stay home from work or school?

Yes. Any person diagnosed with measles should stay away from school or work for 4 days after the rash begins.

### How can you keep from getting it?

The best way to keep from getting measles is to get vaccinated.

Pregnant women and people with a life threatening allergic reaction to gelatin, neomycin, or a previous dose of MMR vaccine should not receive the MMR vaccine. Persons with immune system problems or who are moderately or severely ill should consult with their doctor about whether they should be vaccinated.

For more information, see the CDC's website at [http://www.cdc.gov/ncidod/diseases/submenu/sub\\_measles.htm](http://www.cdc.gov/ncidod/diseases/submenu/sub_measles.htm)