



Hawaii State Department of Health

Disease Investigation Branch

Oahu 586-4586, Maui 984-8213, Hilo 933-0912,

Kona 322-4877, Kauai 241-3563

Tularemia

(Rabbit fever, Deerfly fever)

What is Tularemia?

Tularemia is a bacterial disease of humans and animals caused by the bacteria *Francisella tularensis*. Although many types of household, farm, and wild animals (especially rabbits, hares, muskrats and beavers) can be infected, tularemia is most often spread to people from infected wild rabbits. Tularemia is not known to be present in Hawaii. The bacteria occurs naturally worldwide and could also be used as a bioterrorist weapon.

How do you get it?

People can become infected with tularemia by direct contact (through mucous membranes of the eyes, nose and mouth, or cuts in the skin) with infected animals, inhaling the bacteria from contaminated environments, eating or drinking contaminated food or water or by the bite from an insect vector such as a tick or deerfly. Tularemia is not spread from person to person.

The bacteria can live for weeks to months in soil, water and vegetation.

What are the symptoms of tularemia?

The symptoms of tularemia vary by the route of infection. Initial signs are often flu-like including fever, chills, nausea, headache and joint pain. When tularemia is spread through a scratch or bite, a sore may appear at the site where the bacteria entered the body and lymph nodes near the site may be swollen and painful. Swallowing the bacteria in food or water may cause sore throat, stomach pain, diarrhea and vomiting. Inhaling the bacteria can produce fever, coughing, chest pain, shortness of breath and a pneumonia like illness.

When do symptoms start?

Signs of illness usually begin 3-5 days after exposure to the bacteria, but the onset may vary from 2 to 10 days.

What is the treatment for tularemia?

Antibiotics and other medications can be prescribed by physicians to treat tularemia. Let your physician know if you have had contact with wildlife or the environment in an area where tularemia has been diagnosed.

How can you keep from getting it?

- Avoid contact with wildlife, especially if they appear ill. If you must handle wildlife, wear gloves and wash your hands after touching any animal.
- Wear protective gear (rubber gloves, goggles) when skinning or handling wild animals such as rabbits in areas where tularemia is found.
- Thoroughly cook wild game meats before eating. Wash fruits and vegetables before eating.
- Wear insect repellents and protective clothing to prevent bites of flies, ticks and mosquitoes in areas where tularemia is reported.
- Only drink water and swim at safe water sources.

For more information:

Centers for Disease Control and Prevention:

<http://www.bt.cdc.gov/agent/tularemia/>

Iowa State University Center for Food Security and Public Health:

http://www.cfsph.iastate.edu/FastFacts/pdfs/tularemia_F.pdf